



Assessing neighborhood disorder: Validation of a three-factor observational scale



Miriam Marco^{a,*}, Enrique Gracia^a, José M. Tomás^b, Antonio López-Quílez^c

^a Department of Social Psychology, University of Valencia, Spain

^b Department of Methodology of the Behavioral Sciences, University of Valencia, Spain

^c Department of Statistics and Operations Research, University of Valencia, Spain

ARTICLE INFO

Article history:

Received 20 May 2015

Accepted 25 May 2015

Available online 19 June 2015

Keywords:

Confirmatory factor analysis

Neighborhood disorder

Observational scale

Physical decay

Physical disorder

Spatial clustering

Social disorder

Social disorganization

ABSTRACT

This study presents data on the development and preliminary validation of an observational scale assessing neighborhood disorder. Independent observations by trained raters of neighborhood disorder were conducted in 552 census block groups in the city of Valencia (Spain). Intraclass correlation coefficients assessing inter-rater reliability indicated fair to substantial levels of agreement among raters. Confirmatory factor analyses supported a final three-factor model scale measuring physical disorder, social disorder, and physical decay. Results for the internal consistency showed large composite reliability indices indicating good reliability for all neighborhood disorder factors. Evidence of criterion-related validity was found by exploring associations between neighborhood disorder factors and three neighborhood characteristics: neighborhood socioeconomic status, immigrant concentration, and residential instability. Also for criterion-related validity, Moran's I test results for spatial correlation showed that the three types of neighborhood disorder tend to cluster in space and are not randomly distributed across the city. In general, this paper provides evidence of a reliable and valid observational measure to assess neighborhood disorder.

© 2015 Colegio Oficial de Psicólogos de Madrid. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Evaluación del desorden en los vecindarios: validación de una escala observacional de tres factores

RESUMEN

Este estudio presenta el desarrollo y validación preliminar de una escala observacional para la evaluación del desorden en los vecindarios. Se realizaron observaciones independientes del desorden por evaluadores entrenados en 552 sectores censales de la ciudad de Valencia (España). Los coeficientes de correlación intraclass para la evaluación de la fiabilidad interjueces indicaron unos niveles adecuados de acuerdo entre jueces. Los resultados del análisis factorial confirmatorio apoyaron un modelo final de tres factores: desorden físico, desorden social y deterioro físico. La evaluación de la consistencia interna mediante *composite reliability indices* mostró valores elevados para todos los factores. La validez de criterio fue determinada mediante la exploración de las asociaciones entre los factores de desorden del vecindario y tres características del mismo: estatus socioeconómico, concentración de inmigrantes e inestabilidad residencial. Además, como medida de validez de criterio, el test de Moran que evalúa la correlación espacial mostró que los tres tipos de desorden tienden a agruparse espacialmente y no se distribuyen aleatoriamente en la ciudad. En general, este artículo proporciona evidencias de la fiabilidad y validez de una escala para la medida del desorden en los vecindarios.

© 2015 Colegio Oficial de Psicólogos de Madrid. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Palabras clave:

Análisis factorial confirmatorio

Desorden del vecindario

Deterioro físico

Desorden físico

Distribución especial

Desorden social

Desorganización social

Escala observacional

* Corresponding author.

E-mail address: Miriam.Marco-Francisco@uv.es (M. Marco).

2010; Kawachi & Berkman, 2003; O'Campo et al., 2015; Sampson, 2012; Sampson, Raudenbush, & Earls, 1997). Among these neighborhood characteristics, the concept of neighborhood disorder has played a central role and has received the attention of scholars from different disciplines like sociology, criminology, social psychology or epidemiology. Neighborhood disorder can be defined as "observed or perceived physical and social features of neighborhoods that may signal the breakdown of order and social control, and that can undermine the quality of life" (Gracia, 2014, p. 4325). Examples of neighborhood disorder may include behaviors such as prostitution, drug dealing, and fighting in the streets, or physical characteristics such as abandoned cars, vandalized buildings, or litter in the streets (Sampson & Raudenbush, 1999; Skogan, 1990; Taylor, 2001; Wilson & Kelling, 1982).

The concept of neighborhood disorder can be linked to social disorganization theories and their idea that structural characteristics of neighborhoods, like concentrated disadvantage, can undermine social control and increase levels of violence, crime, and other negative outcomes (Gracia, 2014; Kingston, Huiziga, & Elliot, 2009; Kubrin & Weitzer, 2003; Maimon & Browning, 2010; Park, Burgess, & McKenzie, 1925; Sampson et al., 1997; Shaw & McKay, 1942; Wilson, 1987). Also, the Broken Windows Theory of urban decay has been of particular relevance for the wide appeal of the concept of neighborhood disorder (Wilson & Kelling, 1982). According to this perspective, physical and social cues of neighborhood disorder signal the breakdown of formal and informal social controls leading to further disorder and crime (Gracia, 2014; Perkins, Meeks, & Taylor, 1992; Sampson & Raudenbush, 1999; Skogan, 1990; Taylor, 1997, 2005; Toet & van Schaik, 2012; Wei, Hipwell, Pardini, Beyers, & Loeber, 2005; York Cornwell & Cagney, 2014). According to Gracia (2014) "as neither residents nor external agencies (e.g., police and other authorities) are able or willing to intervene and maintain social order, more disorder is facilitated, and criminal activity is attracted" (p. 4325). Neighborhood disorder would also trigger a number of community processes like fear, insecurity, powerlessness, or mistrust that lead residents to disinvest in and withdraw from community life, increasing social disorganization and neighborhood decline (Geis & Ross, 1998; Kawachi, Kennedy, & Wilkinson, 1999; Kim & Conley, 2011; Ross, Mirowsky, & Pribesh, 2001; Skogan, 1986, 1990). In this regard, neighborhood disorder has been linked to urban decay, concentration of social problems, racial or ethnic segregation, social integration, confidence in the police, or public social control strategies like reporting crime (Gracia, Garcia, & Musitu, 1995; Gracia & Herrero, 2006a, 2006b, 2007; Perkins et al., 1992; Perkins & Taylor, 1996; Ross & Mirowsky, 1999; Skogan, 1990; Taylor, 1997; Toet & van Schaik, 2012).

Although neighborhood disorder has traditionally been studied in relation to street-level outcomes, an increasing body of literature has also examined its influence on processes and outcomes that occur "behind closed doors" (Wright & Benson, 2011), such as parental socialization practices (Gracia, Fuentes, García, & Lila, 2012; Lila & Gracia, 2005; McDonell, 2007; Roosa et al., 2005; Tendulkar, Buka, Dunn, Subramanian, & Koenen, 2010; White, Roosa, Weaver, & Nair, 2009; Worton et al., 2014), child maltreatment (Coulton, Crampton, Irwin, Spilsbury, & Korbin, 2007; Coulton, Korbin, & Su, 1999; Freisthler, Bruce, & Needell, 2007; Freisthler, Merritt, & LaScala, 2006; Garbarino & Sherman, 1980; Gracia & Musitu, 2003; Lila & Gracia, 2005; Martin-Storey et al., 2012), or intimate partner violence (Cunradi, 2007, 2009; Gracia, Herrero, Lila, & Fuente, 2009; Gracia, López-Quílez, Marco, Lladosa, & Lila, 2014, 2015; Kirst, Lazgare, Zhang, & O'Campo, 2015; see Beyer, Wallis, & Hamberger, 2015; Pinchevsky & Wright, 2012, for reviews).

More recently, research on social disorder has also examined its influences on individual well-being indicators like subjective well-being, psychological distress, anxiety, or depression

(García-Ramírez, Balcázar, & de Freitas, 2014; Herrero, Gracia, Fuente, & Lila, 2012; Hill & Angel, 2005; Hombrados-Mendieta & López-Espigares, 2014; Latkin & Curry, 2003; Latkin, German, Hua, & Curry, 2009; O'Campo et al., 2015; Ross & Mirowsky, 2009), and how this may affect negative health behaviors such as low physical activity, heavy drinking, smoking, or obesity (Burdette & Hill, 2008; Echeverría, Diez-Roux, Shea, Borrell, & Jackson, 2008; Hill, Ross, & Angel, 2005; Keyes et al., 2012; O'Campo et al., 2015; Ross & Mirowsky, 2001). Research has also examined the association between neighborhood disorder and different public health issues such as health service usage, low body weight at birth in children, injuries, sexually transmitted diseases, loss of physical function in older adults, and mortality risk (Balfour & Kaplan, 2002; Cohen et al., 2000, 2003; Martin-Storey et al., 2012; Pearl, Braveman, & Abrams, 2001; Winkleby & Cubbin, 2003).

Assessing Neighborhood Disorder

Assessment of neighborhood disorder typically considers two types of disorder, physical and social (Robinson, Lawton, Taylor, & Perkins, 2003; Skogan & Maxfield, 1981; Taylor & Shumaker, 1990). Physical disorder refer to urban landscapes with high levels of decay and deterioration. For example, abandoned houses, graffiti, trash on the streets, abandoned cars, used needles, and vacant lots would exemplify physical disorder (Brunton-Smith, 2011; Garvin, Cannuscio & Branas, 2013; Robinson et al., 2003; Sampson & Raudenbush, 1999; Skogan, 1990; Taylor, 2001, Toet & van Schaik, 2012). Some scholars, however, make a further distinction between physical disorder and physical decay: physical disorder would refer to features like dirt in the streets (litter, bottles, condoms), graffiti, abandoned cars, etc. (i.e., behavioral manifestations), whereas physical decay would refer to structural characteristics that can arise from lack of institutional investments and have long term effects, such as abandoned buildings, burn-out houses, badly deteriorated recreational facilities, etc. (Sampson, 2009; Sampson & Raudenbush, 2004). As Sampson and Raudenbush (2004) argue, it is important to make this distinction because physical disorder is "limited to behavioral manifestations (e.g., graffiti, garbage in the streets) that can be conceptually decoupled from structural resources" (p. 326). Social disorder refer, on the other hand, to events in public places seen as potentially threatening, and can be exemplified by the presence of people taking drugs or alcohol in the street, drug dealing, fights and arguments, presence of homeless people, public drunkenness, street prostitution, high levels of police activity, and other criminal or not criminal activities that create a sense of danger (Gracia, 2014; Gracia & Herrero, 2007; Robinson et al., 2003; Ross & Mirowsky, 2001; Sampson, 2009; Sampson & Raudenbush, 2004). Despite some studies suggesting that physical and social disorder may overlap, being order and disorder two ends of a single continuum (Ross & Mirowsky, 1999; Xu, Fielder, & Flaming, 2005), most studies support the distinction between physical and social disorder (Brunton-Smith, Sindall, & Tarling, 2010; LaGrange, Ferraro, & Supancic, 1992; Sampson & Raudenbush, 2004; Taylor & Shumaker, 1990).

In order to assess neighborhood disorder, researchers generally use three different approaches (McDonell & Waters, 2011; Mooney et al., 2014). One approach, based on a more objective perspective, draws from neighborhood information from governmental or commercial data sources (Cerdá et al., 2009; McDonell, 2007; Mooney et al., 2014). Although these data is freer from the variability and subjectivity of subjective perceptions of disorder (Kubrin, 2008), however, this information is "often collected for administrative purposes, may not fully capture the construct of research interest, and may be collected at a spatial resolution that is not optimal for research purposes" (Mooney et al., 2014, p. 626–627). A second, and widely used, approach is based on resident's perceptions of

their neighborhood physical or social characteristics. A number of limitations have been noted, however, regarding this approach, including “same source bias” (e.g., same source reporting perceived neighborhood disorder and related outcomes), confusion with other psychological constructs (e.g., fear of crime), or the influence of stereotypes and neighborhood prejudices (e.g., racial, ethnic, or socioeconomic composition) in perceptions of disorder (Caughy, O’Campo, & Patterson, 2001; Duncan & Raudenbush, 1999; Gómez, Johnson, Selva, & Sallis, 2004; Mooney et al., 2014; Sampson, 2009; Sampson & Raudenbush, 1999, 2004; Schaefer-McDaniel, Caughy, O’Campo, & Gearey, 2010). Finally, a third approach, that aims to overcome the above limitations, emphasizes the importance of using direct and systematic observations of neighborhood characteristics by trained researchers (Franzini, Caughy, Nettles, & O’Campo, 2008; McDonell, 2007; McDonell & Waters, 2011; Neil, Parke, & McDowell, 2001; Raudenbush & Sampson, 1999; Reiss, 1971; Sampson & Raudenbush, 1999). This approach aims to obtain objective measures of neighborhood conditions, to capture a wide range of factors, which are not always available otherwise, and to allow its replication in other contexts (Caughy et al., 2001; Cohen et al., 2000; Franzini et al., 2008; McDonell & Waters, 2011; Sampson & Raudenbush, 2004; Taylor, 2001).

The Present Study

This study aims to add to this body of research by validating an observational measure of neighborhood disorder in the context of a European city that may differ from the culture and structure of Anglo-Saxon cities where most of this type of measures have been developed and validated (Le Galès & Zagrodzki, 2006; Summers, Cheshire, & Senn, 1999). The availability of a reliable and valid measure of neighborhood disorder in this context may be an important addition to the growing body of literature exploring neighborhood effects (Sampson, 2012). To this end, independent observations by trained raters of neighborhood disorder will be conducted using small-areas of the city as the ecological units (i.e., census block groups, which are the smallest administrative sections of the city available). By using the smallest possible geographical units of the city we expect greater homogeneity of neighborhood characteristics (Ocaña-Riola et al., 2008). Also, by using all census block groups of the city we will obtain greater variability and, as neighborhood characteristics tend to cluster in space, this will provide the possibility to explore the clustering of these characteristics for validation purposes (Gracia et al., 2014).

This study presents data on the development and preliminary validation of an observational instrument to assess neighborhood disorder. The specific objectives of the study are: (1) to test the inter-rater reliability of the scale; (2) to test the factorial validity of the scale using confirmatory factor analysis: we expect a factorial structure reflecting three theoretically a priori factors, physical disorder, social disorder, and physical decay (Sampson & Raudenbush, 2004); (3) to test the reliability of the scale by means of the composite reliability index; and (4) to test the criterion-related validity of the scale also employing structural equations. Drawing from social disorganization theory we expect associations between neighborhood disorder and three neighborhood structural characteristics, central in this theoretical perspective: neighborhood socioeconomic status, immigrant concentration, and residential instability (Caughy et al., 2001; Jones, Pebley, & Sastry, 2011; Kubrin & Weitzer, 2003; McDonell & Waters, 2011; Mooney et al., 2014; Sampson et al., 1997; Sampson & Raudenbush, 1999). Also, for validation purposes, a spatial perspective will be applied. As we expect that disordered neighborhoods will cluster in space, rather than be randomly distributed in the city, spatial correlation analyses will be conducted to test whether neighborhood disorder shows a significant spatial pattern (Bruinsma, Pauwels, Weerman, &

Bernasco, 2013; Gracia et al., 2015; Quick, 2013; Veysey & Messner, 1999).

Method

Sample

This study was conducted in the city of Valencia, the third largest city of Spain. As proxy of neighborhood units we used census block groups that were the smallest administrative unit of the city available. Census block groups can be defined as walkable areas within a few number of city blocks and as they are smaller than census tracks, they are particularly appropriate for neighborhood studies (Gracia et al., 2014; Sampson & Raudenbush, 2004). Observations by trained raters were conducted in each of the 552 census block groups in which the city is divided. The total population for these census block groups was 736,580 inhabitants (2013 data), with an average of 1,334 inhabitants per census block group (ranging from 630 to 2,845).

Measures

Neighborhood disorder observation scale. A neighborhood disorder scale was initially constructed with a total of 20 items based on three dimensions of neighborhood disorder proposed by Sampson and Raudenbush (Sampson & Raudenbush, 2004). Thus, the scale included three theoretically motivated subscales measuring physical disorder, social disorder, and physical decay. *Physical disorder* was defined by 8 items: cigarettes in the street, trash in the street, empty bottles in the street, graffiti, abandoned cars, used condoms and syringes in the street, and political or protest message graffiti. *Social disorder* was defined by 7 items: adults or young people loitering, people drinking alcohol in public, gangs, public intoxication, adults fighting or arguing, selling drugs, and street prostitution. *Physical decay* was defined by 5 items: vacant or abandoned houses, abandoned commercial buildings, vandalized and run-down buildings, deteriorated residential units and deteriorated recreation places. The observations are rated on a 5-point response scale (from 0 = *no presence*, to 4 = *highly present*). Two trained raters walked each census block group in order to complete the observational scale. All observations were made during business hours.

Criterion-related validity measures. Drawing from social disorganization theory, to test criterion-related validity we will explore relationships between neighborhood disorder and three neighborhood characteristics measured at census block group level: neighborhood socioeconomic status, immigrant concentration, and residential instability. The City of Valencia Statistics Office provided these data for each census block group. Neighborhood socioeconomic status was measured with an indicator created through factor analysis (this factor included educational level, property value, percentage of high-end cars, and financial and commercial activities). Immigrant concentration was the percentage of immigrant population in each census block, and residential instability was the proportion of the population who had moved into or out of each census block group during the previous year (rate per 1,000 inhabitants).

Statistical Analysis

To measure inter-rater reliability, two pairs of trained undergraduate students walked a random sample of the census block groups. They observed a subset of 15% of them approximately ($N = 86$). Inter-rater reliability scores were computed per each of the three scales by calculating intraclass correlation coefficients (ICC) due to the quantitative nature of data. This analysis was performed with SPSS 22 for Windows.

Several competing confirmatory factor analyses (CFA) were specified, estimated and tested in Mplus 7.3. According to the ordinal nature of the data and its non-normality, WLSMV (weighted least squares mean and variance corrected) estimation was used, the one recommended in the literature (Finney & DiStefano, 2006). Several criteria were used to assess goodness-of-fit: (a) the chi-square statistic, (b) the comparative fit index (CFI), and (c) the root mean squared error of approximation (RMSEA). A model with a CFI of .95 or larger and a RMSEA of .08 or lower would be indicative of very good fit between the hypothesized model and the data (Hu & Bentler, 1999). Nevertheless, overall fit must be accompanied by a careful diagnosis of the analytical fit (parameter estimates) in the model in order to not blindly use the aforementioned thresholds (Kline, 2011). For model comparison, a modeling approach that uses practical fit indices to determine the overall adequacy of a fitted model has been used as recommended by Cheung and Rensvold (2002) or Little (1997). From this point of view, if a parsimonious model evidences adequate levels of practical fit, then it is preferred over the more complex model. Usually, CFI differences (Δ CFI) are used to evaluate measurement invariance. CFI differences lower than .01 (Cheung & Rensvold, 2002) or .05 (Little, 1997) are usually employed as cut-off criteria.

Additionally, internal consistency of the dimensions in the scale has been estimated with the composite reliability index (CRI). Although Cronbach's coefficient alpha is the most widely used estimator of internal consistency, it has been criticized as being only completely appropriate with essentially tau-equivalent items (and tests) and also by being a lower bound for the true reliability (Raykov, 2004). More explicitly, a tau-equivalent test assumes all items measure the same latent variable, on the same scale, with the same degree of precision, with all true scores being equal (Graham, 2006). When tau-equivalence does not hold, alpha will over- or under-estimate (more often the latter) the population value. An alternative to the coefficient alpha is the CRI, which is usually calculated using estimates from confirmatory factor analyses (Graham, 2006). Accordingly, the more adequate CRI was employed.

Criterion-related validity was established by correlating neighborhood disorder factors with other neighborhood constructs theoretically linked in the literature (Gracia, 2014; Gracia et al., 2015; Kubrin & Weitzer, 2003; Sampson & Raudenbush, 1999; Sampson et al., 1997; Shaw & McKay, 1942). This correlation was obtained within the context of a structural equation modeling in order to prevent as much as possible the correlation attenuation due to measurement error. Specifically, neighborhood disorder was correlated to neighborhood socioeconomic status, immigrant concentration, and residential instability (see Measures section).

To test criterion-related validity we also used a spatial methodology approach. To assess spatial autocorrelation, we computed Moran's I (Moran, 1950) per each of the three subscales or factors, considering as the observation the midpoint of each of the census block groups. We expected a significant spatial distribution of neighborhood disorder, rather than a random distribution, because we expect that disorder, as other neighborhood characteristics, will show a tendency towards spatial clustering (Bruinsma et al., 2013; DiMaggio, 2015; Quick, 2013; Veysey & Messner, 1999).

Results

Inter-Rater Reliability

Intraclass correlation coefficients were computed to assess inter-rater agreement for the three subscales (see Odgers, Caspi, Bates, Sampson, & Moffitt, 2012, for a similar approach). Intraclass correlations ranged from .25 to .71 (see Table 1). Landis and Koch's (1977) criteria were used to interpret results regarding inter-rater

agreements: < .20 slight, .21 – .40 fair, .41 – .60 moderate, .61 – .80 substantial, and .81 – 1 almost perfect agreement. Our results indicated fair to substantial levels of agreement between raters (Landis & Koch, 1977). Social disorder obtained the lowest value and physical disorder and physical decay showed similar results.

Table 1
Inter-rater Agreement. Intra-class Correlations Coefficients (ICC)

Scales	No. of items	M (SD)	ICC ₁	ICC ₂
Physical disorder	8	5.97 (3.54)	.55***	.71***
Social disorder	7	0.57 (0.79)	.25***	.40***
Physical decay	5	2.64 (2.82)	.46***	.63***

Note. ICC₁ = index of reliability for a single rater.

ICC₂ = index of reliability for multiple raters averaged together.

* < .05, ** < .01, *** < .001

Confirmatory Factor Analysis

Three a priori competing models were specified. The theoretical model that supports the content validity of the scale a priori hypothesizes three dimensions: physical disorder, social disorder, and physical decay. Indicators were developed to tap these three theoretical dimensions. However, there was doubt about whether two of these dimensions could be too overlapped to have discriminant validity: physical disorder and physical decay. Accordingly, another a priori model was specified with two dimensions: social disorder and all the indicators of physical disorder and physical decay specified to load onto a single dimension. Finally, the most parsimonious latent structure, a one-factor model underlying all the indicators, was also specified. Goodness-of-fit indices for these three a priori models are shown in Table 2. Model fit for the three models was extremely poor, and none of them could be retained as a good approximation to the observed data. Nevertheless, the three-factor model showed a *relative* better fit compared to the other two models: Δ CFI = .06, compared to the one-factor model and Δ CFI = .04, compared to the two-factor model. Taking this information into account, plus the fact that the theoretical model that supports the scale was three-dimensional, this model was retained for further psychometric scrutiny.

Table 2
Goodness of Fit Indices for the Tested Models

	χ^2	df	p	CFI	RMSEA	90% CI
One-factor model	1471.75	152	< .001	.643	.12	.12 - .13
Two-factor model	1407.56	151	< .001	.660	.12	.11 - .12
Three-factor model	1272.59	70	< .001	.700	.11	.11 - .12
Modified three-factor model	278.61	87	< .001	.940	.06	.05 - .07

A careful look at the factor loadings, together with the lack of variability in some indicators, allows to remove some of them. Those removed lacked variability and/or had poor consistency with their dimension. The final version of the questionnaire was presumed to measure three factors (physical disorder, social disorder, and physical decay) with 5, 6, and 4 indicators each (see Appendix). This depurated version of the original scale was tested and its goodness-of-fit indices are shown in Table 2. There was a huge improvement in model fit, and it can be said that the model seems to adequately represent the observed data. Factor loadings are shown in Figure 1. They were all statistically significant ($p < .01$) and, in general, pretty large. These results are indicative of good analytical fit.

Internal Consistency

Reliability (internal consistency) estimates were calculated for each dimension or factor in the scale. The calculated reliability




Figure 1. Standardized factor loadings and correlations for the confirmatory factor analyses of the Neighborhood Disorder Observational Scale.

Note. All coefficients statistically significant ($p < .01$); Intoxicat = intoxication; Dealing = drug dealing; Vacant = vacant houses; Vandal. = vandalized buildings; Residen. = residential deterioration; Deter. = deteriorated facilities.

estimates were composite reliability indices (CRI), as already mentioned in the method section. All the CRI were large, indicating a good reliability for all the dimensions. Specifically, CRI for physical disorder was .83, .82 was the estimate for social disorder, and .82 the internal consistency estimated for physical decay.

Criterion-Related Validity

Finally, evidence of criterion-related validity was also found. Three criteria were used (neighborhood socioeconomic status, immigrant concentration, and residential instability). The relationship between the criteria and the latent variables were calculated at the latent level, not the observed one. A first structural model included the measurement model found to fit well to the data, plus the first criterion (neighborhood socioeconomic status). Again, this structural model fitted the data well, $\chi^2(99) = 339.78$, $p < .001$; CFI = .93, RMSEA = .066 [.059 - .074]. The correlations between neighborhood socioeconomic status and the criteria and the factors were: $\rho = -.13$, $p < .01$ with physical disorder; $\rho = .04$, $p > .05$ with social disorder; and $\rho = -.29$, $p < .01$ with physical decay. With respect to immigrant concentration, the structural model also adequately fitted the data, $\chi^2(99) = 309.38$, $p < .001$; CFI = .93, RMSEA = .062 [.054 - .070]. The correlations among the criterion and the factors were: $\rho = .44$, $p > .05$ with physical disorder; $\rho = .13$, $p < .05$ with social disorder; and $\rho = .21$, $p < .01$ with physical decay. Finally, a third structural model was specified and tested to relate the three factors with residential instability. Again the model fitted the data well, $\chi^2(99) = 333.54$, $p < .001$; CFI = .92, RMSEA = .066 [.058 - .073], and its correlations with the factors were: $\rho = .05$, $p > .05$ with physical disorder; $\rho = .15$, $p < .05$ with social disorder; and $\rho = .22$, $p < .01$ with physical decay.

Also, for criterion-related validity purposes, tests for spatial correlation were conducted for the three dimensions. Results showed spatial autocorrelation in the three scales as all Moran's I values

were positive and significant ($p < .001$). Moran's I values for the three measures were .20 for social disorder, .39 for physical decay, and .49 for physical disorder, indicating a stronger spatial pattern for physical disorder and physical decay than for social disorder. These results indicate a positive non-random distribution of all types of neighborhood disorder in the city (i.e., rather than being randomly distributed across the city, they tend to cluster in space).

Discussion

In this paper we describe the development and the psychometric properties of a preliminary validation of an observational scale assessing neighborhood disorder. This scale was implemented in the city of Valencia (Spain), using independent observations of all census block groups of the city conducted by trained raters. In general, results have shown that this scale is a psychometrically sound and valid instrument to assess three neighborhood disorder dimensions: physical disorder, social disorder, and physical decay.

Results regarding inter-rater reliability showed fair to substantial levels of agreement (Landis & Koch, 1977), with stronger agreements for physical disorder and physical decay factors, and lower inter-rater reliability for social disorder. Although these results are slightly lower than in other studies (e.g., Caughy et al., 2001; Franzini et al., 2008; Jones et al., 2011), they are, however, comparable to others (e.g., Mooney et al., 2014; Wei et al., 2005). As to why social disorder showed lower levels of agreement, one possibility is that social disorder cues such as fights or public intoxication tend to be less stable over time (e.g., depending on the time of the day) than other physical features of neighborhoods, like those indicating physical decay and disorder, that are more temporally stable (Jones et al., 2011).

With respect to the factor structure of the observational scale, our analyses aimed to validate a three theoretically-based neighborhood disorder dimensions (Sampson & Raudenbush, 2004). To

this end, several competing confirmatory factor analyses were first estimated and discarded, yielding a final and depurated version of the scale with good analytical fit. This final scale supported a three-factor model measuring physical disorder, social disorder, and physical decay, as theorized by Sampson and Raudenbush (2004).

This final three-factor model was obtained after removing some items for their low variability or poor consistency with their dimension. Although these items are usually present in other scales measuring neighborhood disorder, however, indicators such as abandoned cars, used condoms and syringes (physical disorder), prostitution (social disorder), or abandoned commercial buildings (physical decay) were discarded in the final model. The presence of these items was very low and, in the case of syringes, there was no presence at all in any of the census block groups observed. Although the time of the observation may have influenced the low presence of some of these indicators (e.g., prostitution), another possible explanation is that, given the characteristics of cities like Valencia, with high density in a relative small area (Le Galès & Zagrodzki, 2006; Summers et al., 1999), some of these indicators could be more present in the outskirts of the city, and therefore outside of the boundaries where the observations were made. This also suggests that differences between the present scale and others developed elsewhere may reflect context-specific features of the cities. Given that these results may be context-dependent, we do not favour the uncritical use of the short-version of the originally proposed scale distilled for this particular study. On the contrary, careful theoretical considerations previous to the use of this scale in other cities should consider whether some (or all) of the items removed could have enough variability and importance as to be included in the instrument. Clearly, a subsequent depuration of the scale according to its psychometric properties is always possible. On the other hand, results for internal consistency of the final three-factor scale by confirmatory analyses also supported its reliability, with CRI values between .82 and .83 for the three scales.

Two different criterion-related validity tests of this observational measure of neighborhood disorder were conducted. For the first one, and drawing from social disorganization theory, we explored associations with three criteria tapping neighborhood characteristics central to this theoretical perspective: neighborhood socioeconomic status, immigrant concentration, and residential instability, measured at the census block group level. As expected, correlations between measures of neighborhood disorder and these structural characteristics of neighborhoods were mostly in the expected direction (Kubrin & Weitzer, 2003; Sampson & Raudenbush, 1999; Sampson et al., 1997), although with stronger associations for physical decay. For example, high levels of physical decay were significantly associated with lower neighborhood socioeconomic status, higher rates of immigrant concentration, and higher residential instability. Social disorder was positively related to levels of immigrant concentration and residential instability, but its association with neighborhood socioeconomic status was not statistically significant. Physical disorder was also negatively related to neighborhood socioeconomic status; however, associations with immigrant concentration and residential instability did not reach significance. These results partly support previous research where significant associations between disorder and a number of neighborhood characteristics were also found, especially those regarding the relationship between physical disorder and decay and neighborhood socioeconomic indicators (Caughy et al., 2001; Jones et al., 2011; McDonnell & Waters, 2011; Mooney et al., 2014; Sampson & Raudenbush, 1999). Also, some of our results support studies that fail to find a significant relationship between physical disorder and residential instability (Sampson & Raudenbush, 1999), or between physical disorder and immigrant concentration (Jones et al., 2011). It is interesting to note that physical decay was

associated with all neighborhood characteristics (socioeconomic status, immigrant concentration, and residential instability), suggesting that differentiating between physical disorder and physical decay is an important theoretical distinction that may provide a more detailed analysis when assessing neighborhood disorder and exploring its relationships with different outcomes and processes (Sampson, 2009; Sampson & Raudenbush, 2004).

As we expected that disordered neighborhoods would tend to cluster together in space (Bruinsma et al., 2013; DiMaggio, 2015; Quick, 2013; Veysey & Messner, 1999), for the second criterion-related validity test we used a spatial analytical approach to assess the spatial distribution of the different types of neighborhood disorder across all census block groups observed. Results showed that the three types of disorder (physical disorder, social disorder, and physical decay) were spatially clustered, confirming that they were not randomly distributed in the city. This reflects the existence of different areas of the cities where neighborhood disorder tends to concentrate and shows that this neighborhood risk factor tend to cluster in space. These results support the idea that different manifestations of neighborhood disorder, as other characteristics of the cities, are not randomly distributed in space. As illustrated by a growing body of literature linking neighborhood disorder with a wide array of outcomes, including crime, violence, or health, the spatially patterned nature of this risk factor makes more likely that related outcomes will also be spatially patterned (Cunradi, Mair, Ponicki, & Remer, 2011; Diez-Roux & Mair, 2010; Freisthler et al., 2007; Gracia et al., 2015; Kawachi & Berkman, 2003; Law, Quick, & Chan, 2014; O'Campo et al., 2015; Sampson, 2012).

Finally, this study has both strengths and limitations. Among the strengths, the use of independent observations of neighborhood conditions, rather than residents' subjective perceptions, allows to overcome some of the limitations noted in the literature regarding this later approach (Caughy et al., 2001; Mooney et al., 2014; Sampson & Raudenbush, 1999; Schaefer-McDaniel et al., 2010). Relatedly, for the observation of neighborhood disorder at the level of ecological units (rather than the personal level), we also use a high-resolution approach. We used the smallest administrative units available (i.e., census block groups) that allow greater homogeneity and precision than lower resolutions such as census tracts or postal codes, commonly used in other neighborhood studies (Beyer et al., 2015; Bursik, Grasmick, & Chamlin, 1990; Kaufman, Dole, Savitz, & Herring, 2003; Campo, Xue, Wang, & Caughy, 1997). By using small-area units, we also reduced potential ecological bias, as this resolution is closer to the individual level (Gracia et al., 2015; Lawson, 2006; Ocaña-Riola et al., 2008). We used census block groups in our study, which substantially reduced this potential bias. Finally, we used all census block groups of the city, rather than selecting only a sample of them, which provided greater variability and the possibility to explore potential significant spatial patterns in the distribution of neighborhood disorder across the city (Caughy et al., 2001; Mooney et al., 2014). In this regard, the use of spatial methods to complement the criterion-related validity of our observational scale is an important addition to the existing literature, as neighborhoods, from this perspective, are not treated as independent units (Gorman, Gruenewald, & Waller, 2013; Mooney et al., 2014; Morenoff, Sampson, & Raudenbush, 2001). Although widely used in epidemiological studies (Lawson, 2009), and despite its advantages, this methodological approach is still uncommon in neighborhood studies, and future research would clearly benefit from incorporating a spatial perspective (Cunradi et al., 2011; Gracia et al., 2014, 2015; Law & Quick, 2013; Law et al., 2014; Sparks, 2011). As for limitations, as noted above, some measures of neighborhood disorder may have been affected by the time of the day they were observed. Our observations were limited to business hours, and the same places may have shown different characteristics at night (Caughy et al., 2001). Future

research would benefit from including different observations during the day, and revisiting the same areas during night hours. Also, other neighborhood indicators such as trash in the streets may be present only at specific moments, and repeated observations of the same area would be advisable, although clearly more costly (Wei et al., 2005). In this regard, recently new technologies, such as virtual environments, or Google Street View, provide powerful and easy accessible tools that may help to enhance neighborhood research (Odgers et al., 2012; Toet & van Schaik, 2012).

In conclusion, this paper provides evidence of a reliable and valid observational measure to assess neighborhood disorder. Adequate measures to assess neighborhood characteristics are important research and intervention tools, as they are key to better understanding neighborhood processes, as well as to evaluate related outcomes and monitor changes after grass-roots efforts or official initiatives to reduce neighborhood inequalities.

Conflict of Interest

The authors of this article declare no conflict of interest.

Financial Support

This research was supported by the Spanish Ministerio de Economía y Competitividad (PSI2014-54561-P). Miriam Marco was supported by the FPU program of the Spanish Ministerio de Educación, Cultura y Deporte (FPU2013/00164).

Appendix. Neighborhood Disorder Observation Scale

Item no.	Scale	Item content *
1	Physical disorder	Colillas en la calle / Cigarettes in the street
2	Physical disorder	Basura en la calle o acera / Trash in the street
3	Physical disorder	Botellas o latas vacías de cerveza u otras bebidas en la calle / Empty bottles or cans in the street
4	Physical disorder	Graffitis / Graffiti
5	Physical disorder	Pintadas de carácter político o reivindicativo / Political or protest message graffiti
1	Social disorder	Jóvenes o adultos merodeando por el barrio / Adults or young people loitering
2	Social disorder	Gente bebiendo alcohol en la vía pública / People drinking alcohol in public
3	Social disorder	Bandas / Gangs
4	Social disorder	Gente borracha o drogada por la calle / Public intoxication
5	Social disorder	Peleas o discusiones agresivas entre jóvenes o adultos / Adults or young people fighting or arguing
6	Social disorder	Venta de droga / Selling drugs
1	Physical decay	Casas vacías / Vacant houses
2	Physical decay	Viviendas abandonadas, quemadas o tapiadas / Abandoned, vandalized and run-down buildings
3	Physical decay	Zonas residenciales muy deterioradas / Deteriorated residential units
4	Physical decay	Zonas recreativas muy deterioradas / Deteriorated recreation places

Note. * Items in Spanish in the original scale / item translation for information purposes

References

- Balfour, J. L., & Kaplan, G. A. (2002). Neighborhood environment and loss of physical function in older adults: Evidence from the Alameda County study. *American Journal of Epidemiology*, 155, 507–551. <http://dx.doi.org/10.1093/aje/k155.6.507>
- Beyer, K., Wallis, A. B., & Hamberger, L. K. (2015). Neighborhood environment and intimate partner violence: A systematic review. *Trauma Violence Abuse*, 16, 16–47. <http://dx.doi.org/10.1177/1524838013515758>
- Bruinsma, G. J. N., Pauwels, L. J. R., Weerman, F. M., & Bernasco, W. (2013). Social disorganization, social capital, collective efficacy and the spatial distribution of crime and offenders: An empirical test of six neighbourhood models for a Dutch city. *British Journal of Criminology*, 53, 942–963. <http://dx.doi.org/10.1093/bjc/azt030>
- Brunton-Smith, I. (2011). Untangling the relationship between fear of crime and perceptions of disorder: Evidence from a longitudinal study of young people in England and Wales. *British Journal of Criminology*, 51, 885–899. <http://dx.doi.org/10.1093/bjc/azr064>
- Brunton-Smith, I., Sindall, K., & Tarling, R. (2010). *The effect of demographic make-up on perceptions of anti-social behaviour in London as measured by the British Crime Survey and 2008 Place Survey*. London: Government Office for London.
- Burdette, A. M., & Hill, T. D. (2008). An examination of processes linking perceived neighborhood disorder and obesity. *Social Science and Medicine*, 67, 38–46. <http://dx.doi.org/10.1016/j.socscimed.2008.03.029>
- Bursik, R. J., Grasmick, H. G., & Chamlin, M. B. (1990). The effects of longitudinal arrest patterns on the development of robbery trends at the neighborhood level. *Criminology*, 28, 431–450. <http://dx.doi.org/10.1111/j.1745-9125.1990.tb01333.x>
- Caughy, M. O., O'Campo, P. J., & Patterson, J. (2001). A brief observational measure for urban neighborhoods. *Health & Place*, 7, 225–236. [http://dx.doi.org/10.1016/S1353-8292\(01\)00012-0](http://dx.doi.org/10.1016/S1353-8292(01)00012-0)
- Cerdá, M., Tracy, M., Messner, S. F., Vlahov, D., Tardiff, K., & Gelea, S. (2009). Misdemeanor policing, physical disorder, and gun-related homicide: A spatial analysis test of "broken-windows" theory. *Epidemiology*, 20, 533–541. <http://dx.doi.org/10.1097/EDE.0b013e3181a48a99>
- Cheung, G. W., & Rensvold, R. B. (2002). Evaluating goodness-of-fit indexes for testing MI. *Structural Equation Modeling*, 9, 235–255.
- Cohen, D. A., Mason, K., Bedimo, A., Scribner, R., Basolo, V., & Farley, T. A. (2003). Neighborhood physical conditions and health. *American Journal of Public Health*, 93, 467–471.
- Cohen, D., Spear, S., Scribner, R., Kissinger, P., Mason, K., & Wildgen, J. (2000). Broken windows" and the risk of gonorrhoea. *American Journal of Public Health*, 90, 230–236. <http://dx.doi.org/10.2105/AJPH.90.2.230>
- Coulton, C. L., Crampton, D. S., Irwin, M., Spilsbury, J. C., & Korbin, J. E. (2007). How neighborhoods influence child maltreatment: A review of the literature and alternative pathways. *Child Abuse & Neglect*, 31, 1117–1142. <http://dx.doi.org/10.1016/j.chiabu.2007.03.023>
- Coulton, C. J., Korbin, J. E., & Su, M. (1999). Neighborhoods and child maltreatment: A multilevel study. *Child Abuse & Neglect*, 23, 1019–1040. [http://dx.doi.org/10.1016/S0145-2134\(99\)00076-9](http://dx.doi.org/10.1016/S0145-2134(99)00076-9)
- Cunradi, C. (2007). Drinking level, neighborhood social disorder, and mutual intimate partner violence. *Alcoholism: Clinical and Experimental Research*, 31, 1012–1019.
- Cunradi, C. B. (2009). Intimate partner violence among Hispanic men and women: The role of drinking, neighborhood disorder, and acculturation-related factors. *Violence Victimization*, 24, 83–97. <http://dx.doi.org/10.1891/0886-6708.24.1.83>
- Cunradi, C. B., Mair, C., Ponicki, W., & Remer, L. (2011). Alcohol outlets, neighborhood characteristics, and intimate partner violence: Ecological analysis of a California city. *Journal of Urban Health*, 88, 191–200. <http://dx.doi.org/10.1007/s11524-011-9549-6>
- Diez-Roux, A. V., & Mair, C. (2010). Neighborhoods and health. *Annals of the New York Academy of Sciences*, 1186, 125–145. <http://dx.doi.org/10.1111/j.1749-6632.2009.05333.x>
- DiMaggio, C. (2015). Small-area spatiotemporal analysis of pedestrian and bicyclist injuries in New York City. *Epidemiology*, 26, 247–254. <http://dx.doi.org/10.1097/EDE.0000000000000222>
- Duncan, G. J., & Raudenbush, S. W. (1999). Assessing the effects of context in studies of children and youth development. *Educational Psychologist*, 34, 29–41. <http://dx.doi.org/10.1207/s15326985ep3401.3>
- Echeverría, S., Diez-Roux, A. V., Shea, S., Borrell, L. N., & Jackson, S. (2008). Associations of neighborhood problems and neighborhood social cohesion with mental health and health behaviors: The multi-ethnic study of atherosclerosis. *Health & Place*, 14, 853–865. <http://dx.doi.org/10.1016/j.healthplace.2008.01.004>
- Finney, S. J., & DiStefano, C. (2006). Non-normal and categorical data in SEM. In G. R. Hancock, & R. O. Mueller (Eds.), *Structural equation modelling: A second course* (pp. 269–314). Greenwich, CO: Information Age Publishing.
- Franzini, L., Caughy, M. O., Nettles, S. M., & O'Campo, P. (2008). Perceptions of disorder: Contributions of neighborhood characteristics to subjective perceptions of disorder. *Journal of Environmental Psychology*, 28, 83–93. <http://dx.doi.org/10.1016/j.jenvp.2007.08.003>
- Freisthler, B., Bruce, E., & Needell, B. (2007). Understanding the geospatial relationship of neighborhood characteristics and rates of maltreatment for black, Hispanic, and white children. *Social Work*, 52, 7–16. <http://dx.doi.org/10.1093/sw/52.1.7>
- Freisthler, B., Merritt, D. H., & LaScala, E. A. (2006). Understanding the ecology of child maltreatment: A review of the literature and directions for future research. *Child Maltreatment*, 11, 263–280. <http://dx.doi.org/10.1177/107759506289524>
- Garbarino, J., & Sherman, D. (1980). High-risk neighborhoods and high-risk families: The human ecology of child maltreatment. *Child Development*, 51, 188–198.
- García-Ramírez, M., Balcázar, F., & de Freitas, C. (2014). Community psychology contributions to the study of social inequalities, well-being and social justice. *Psychosocial Intervention*, 23, 79–81. <http://dx.doi.org/10.1016/j.psi.2014.07.009>
- Garvin, E. C., Cannuscio, C. C., & Branas, C. C. (2013). Greening vacant lots to reduce violent crime: A randomised controlled trial. *Injury Prevention*, 19, 198–203. <http://dx.doi.org/10.1136/injuryprev-2012-040439>

- Geis, K. J., & Ross, C. E. (1998). A new look at urban alienation: The effect of neighborhood disorder on perceived powerlessness. *Social Psychology Quarterly*, 61, 232–246.
- Gómez, J. E., Johnson, B. A., Selva, M., & Sallis, J. F. (2004). Violent crime and outdoor physical activity among inner-city youth. *Preventive Medicine*, 39, 876–881. <http://dx.doi.org/10.1016/j.ypmed.2004.03.019>
- Gorman, D. M., Gruenewald, P. J., & Waller, L. A. (2013). Linking places to problems: Geospatial theories of neighborhoods, alcohol and crime. *Geojournal*, 78, 417–428. <http://dx.doi.org/10.1007/s10708-011-9425-7>
- Gracia, E. (2014). Neighborhood disorder. In A. C. Michalos (Ed.), *Encyclopedia of quality of life and well-being research* (pp. 4325–4328). Dordrecht, Netherlands: Springer.
- Gracia, E., Fuentes, M. C., García, F., & Lila, M. (2012). Perceived neighborhood violence, parenting styles, and developmental outcomes among Spanish adolescents. *Journal of Community Psychology*, 40, 1004–1021. <http://dx.doi.org/10.1002/jcop.21512>
- Gracia, E., García, F., & Musitu, G. (1995). Macrosocial determinants of social integration: Social class and area effect. *Journal of Community and Applied Social Psychology*, 5, 105–119. <http://dx.doi.org/10.1002/casp.2450050204>
- Gracia, E., & Herrero, J. (2006a). La comunidad como fuente de apoyo social: evaluación e implicaciones en los ámbitos individual y comunitario [The community as a source of social support: evaluation and implications at the individual and community levels]. *Revista Latinoamericana de Psicología*, 38, 327–342.
- Gracia, E., & Herrero, J. (2006b). Perceived neighborhood social disorder and resident's attitudes toward reporting child maltreatment. *Child Abuse & Neglect*, 30, 357–365. <http://dx.doi.org/10.1016/j.chiabu.2005.11.001>
- Gracia, E., & Herrero, J. (2007). Perceived neighborhood social disorder and attitudes toward reporting domestic violence against women. *Journal of Interpersonal Violence*, 22, 737–752. <http://dx.doi.org/10.1177/0886260507300755>
- Gracia, E., Herrero, J., Lila, M., & Fuente, A. (2009). Perceived neighborhood social disorder and attitudes toward domestic violence against women in Latin-American immigrants. *The European Journal of Psychology Applied to Legal Context*, 1, 25–43.
- Gracia, E., & Musitu, G. (2003). Social isolation from communities and child maltreatment: A cross-cultural comparison. *Child Abuse & Neglect*, 27, 153–168. [http://dx.doi.org/10.1016/S0145-2134\(02\)00538-0](http://dx.doi.org/10.1016/S0145-2134(02)00538-0)
- Gracia, E., López-Quílez, A., Marco, M., Lladosa, S., & Lila, M. (2014). Exploring neighborhood influences on small-area variations in intimate partner violence risk: A Bayesian random-effects modeling approach. *International Journal of Environmental Research and Public Health*, 11, 866–882. <http://dx.doi.org/10.3390/ijerph110100866>
- Gracia, E., López-Quílez, A., Marco, M., Lladosa, S., & Lila, M. (2015). The spatial epidemiology of intimate partner violence: Do neighborhoods matter? *American Journal of Epidemiology*, <http://dx.doi.org/10.1093/aje/kwv016>
- Graham, J. M. (2006). Congeneric and essentially tau-equivalent estimates of score reliability: What they are and how to use them. *Educational and Psychological Measurement*, 66, 930–944. <http://dx.doi.org/10.1177/0013164406288165>
- Herrero, J., Gracia, E., Fuente, A., & Lila, M. (2012). Desorden social, integración social y bienestar subjetivo en inmigrantes latinoamericanos en España [Social disorder, social integration, and subjective well-being among Latin-American immigrants in Spain]. *Anales de Psicología*, 28, 505–514. <http://dx.doi.org/10.6018/analesps.28.2.148721>
- Hill, T. D., & Angel, R. J. (2005). Neighborhood disorder, psychological distress, and heavy drinking. *Social Science & Medicine*, 61, 965–975. <http://dx.doi.org/10.1016/j.socscimed.2004.12.027>
- Hill, T. D., Ross, C. E., & Angel, R. J. (2005). Neighborhood disorder, psychophysiological distress, and health. *Journal of Health and Social Behavior*, 46, 170–186. <http://dx.doi.org/10.1177/002214650504600204>
- Hombrados-Mendieta, I., & López-Espigares, T. (2014). Dimensiones del sentido de comunidad que predicen la calidad de vida residencial en barrios con diferentes posiciones socioeconómicas [Sense of community: Dimensions predicting residential quality of life in neighborhoods with different socioeconomic status]. *Psychosocial Intervention*, 23, 159–167. <http://dx.doi.org/10.1016/j.psi.2014.08.001>
- Hu, L., & Bentler, P. M. (1999). Cut-off criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling*, 6, 1–55. <http://dx.doi.org/10.1080/10705519909540118>
- Jones, F. M., Pebley, A. R., & Sastry, N. (2011). Eyes on the block: Measuring urban physical disorder through in-person observation. *Social Science Research*, 40, 523–537. <http://dx.doi.org/10.1016/j.ssresearch.2010.11.007>
- Kaufman, J. S., Dole, N., Savitz, D. A., & Herring, A. H. (2003). Modeling community-level effects on preterm birth. *Annals of Epidemiology*, 13, 377–384. [http://dx.doi.org/10.1016/S1047-2797\(02\)00480-5](http://dx.doi.org/10.1016/S1047-2797(02)00480-5)
- Kawachi, I., & Berkman, L. F. (2003). *Neighborhoods and health*. New York, NY: Oxford University Press.
- Kawachi, I., Kennedy, B. P., & Wilkinson, R. G. (1999). Crime: Social disorganization and relative deprivation. *Social Science & Medicine*, 48, 719–731. [http://dx.doi.org/10.1016/S0277-9536\(98\)00400-6](http://dx.doi.org/10.1016/S0277-9536(98)00400-6)
- Keys, K. M., McLaughlin, K. A., Koenen, K. C., Goldmann, E., Uddin, M., & Galea, S. (2012). Child maltreatment increases sensitivity to adverse social contexts: Neighborhood physical disorder and incident binge drinking in Detroit. *Drug and Alcohol Dependence*, 122, 77–85. <http://dx.doi.org/10.1016/j.drugalcdep.2011.09.013>
- Kim, J., & Conley, M. E. (2011). Neighborhood disorder and the sense of personal control: Which factors moderate the association? *Journal of Community Psychology*, 39, 894–907. <http://dx.doi.org/10.1002/jcop.20476>
- Kingston, B., Huizinga, D., & Elliott, D. S. (2009). A test of social disorganization theory in high-risk urban neighborhoods. *Youth & Society*, 41, 53–79. <http://dx.doi.org/10.1177/0044118X09338343>
- Kirst, M., Lazgare, L. P., Zhang, Y. J., & O'Campo, P. (2015). The effects of social capital and neighborhood characteristics on intimate partner violence: A consideration of social resources and risks. *American Journal of Community Psychology*, 55, 314–325. <http://dx.doi.org/10.1007/s10464-015-9716-0>
- Kline, R. B. (2011). *Principles and practice of structural equation modeling*. New York, NY: The Guilford Press.
- Kubrin, C. E. (2008). Making order of disorder: A call for conceptual clarity. *Criminology & Public Policy*, 7, 203–213. <http://dx.doi.org/10.1111/j.1745-9133.2008.00502.x>
- Kubrin, C. E., & Weitzer, R. (2003). New directions in social disorganization theory. *Journal of Research in Crime and Delinquency*, 40, 374–402. <http://dx.doi.org/10.1177/002242780325623>
- LaGrange, R. L., Ferraro, K. F., & Supancic, M. (1992). Perceived risk and fear of crime: Role of social and physical incivilities. *Journal of Research in Crime and Delinquency*, 29, 311–334. <http://dx.doi.org/10.1177/0022427892029003004>
- Landis, J., & Koch, G. (1977). The measurement of observer agreement for categorical data. *Biometrics*, 33, 159–174. <http://dx.doi.org/10.2307/2529310>
- Latkin, C. A., & Curry, A. D. (2003). Stressful neighborhoods and depression: A prospective study of the impact of neighborhood disorder. *Journal of Health and Social Behavior*, 44, 34–44. <http://dx.doi.org/10.2307/1519814>
- Latkin, C. A., German, D., Hua, W., & Curry, A. D. (2009). Individual-level influences on perceptions of neighborhood disorder: A multilevel analysis. *Journal of Community Psychology*, 37, 122–133. <http://dx.doi.org/10.1002/jcop.20284>
- Law, J., & Quick, M. (2013). Exploring links between juvenile offenders and social disorganization at a large map scale: A Bayesian spatial modeling approach. *Journal of Geographical Systems*, 15, 89–113. <http://dx.doi.org/10.1007/s10109-012-0164-1>
- Law, J., Quick, M., & Chan, P. (2014). Bayesian spatio-temporal modeling for analysing local patterns of crime over time at the small-area level. *Journal of quantitative criminology*, 30, 57–78. <http://dx.doi.org/10.1007/s10940-013-9194-1>
- Lawson, A. B. (2006). *Statistical methods in spatial epidemiology* (2nd ed.). New Jersey: John Wiley and Sons.
- Lawson, A. B. (2009). *Bayesian disease mapping: hierarchical modeling in spatial epidemiology*. Boca Raton, FL: CRC Press.
- Le Galès, P., & Zagrodzki, M. (2006). *Cities are back in town: The US/Europe comparison*. Cahier Européen numéro 05/06 du Pôle Ville/métropolis/cosmopolis. Paris: Centre d'Etudes Européennes de Sciences Po. Retrieved from: <http://blogs.sciences-po.fr/recherche-villes/files/2010/01/cahievillie0606.pdf>
- Lila, M., & Gracia, E. (2005). Determinantes de la aceptación-rechazo parental [Determinants of parental acceptance-rejection]. *Psicothema*, 17, 107–111.
- Little, T. D. (1997). Mean and covariance structures (MACS) analyses of cross-cultural data: Practical and theoretical issues. *Multivariate Behavioral Research*, 32, 53–76. http://dx.doi.org/10.1207/s15327906mbr3201_3
- Maimon, D., & Browning, C. R. (2010). Unstructured socializing, collective efficacy, and violent behavior among urban youth. *Criminology*, 48, 443–474. <http://dx.doi.org/10.1111/j.1745-9125.2010.00192.x>
- Martin-Storey, A., Temcheff, C. E., Ruttell, P. L., Serbin, L. A., Stack, D. M., Schwartzman, A. E., & Ledingham, J. E. (2012). Perception of neighborhood disorder and health service usage in a Canadian sample. *Annals of Behavioral Medicine*, 43, 162–172. <http://dx.doi.org/10.1007/s12160-011-9310-0>
- McDonnell, J. R. (2007). Neighborhood characteristics, parenting, and children's safety. *Social Indicators Research*, 83, 177–199. <http://dx.doi.org/10.1007/s1205-006-9063-5>
- McDonnell, J. R., & Waters, T. J. (2011). Construction and validation of an observational scale of neighborhood characteristics. *Social Indicators Research*, 104, 439–457. <http://dx.doi.org/10.1007/s1205-010-9755-8>
- Mooney, S. J., Bader, M. D., Lovasi, G. S., Neckerman, K. M., Teitler, J. O., & Rundle, A. G. (2014). Validity of an ecometric neighborhood physical disorder measure constructed by virtual street audit. *American Journal of Epidemiology*, 180, 626–635. <http://dx.doi.org/10.1093/aje/kwu180>
- Moran, P. A. P. (1950). Notes on continuous stochastic phenomena. *Biometrika*, 37, 17–23.
- Morenoff, J., Sampson, R. J., & Raudenbush, S. (2001). Neighborhood inequality, collective efficacy, and the spatial dynamics of urban violence. *Criminology*, 39, 517–560. <http://dx.doi.org/10.1111/j.1745-9125.2001.tb00932.x>
- O'Campo, P., Wheaton, B., Nisenbaum, R., Glazier, R. H., Dunn, J. R., & Chambers, C. (2015). The neighborhood effects on health and well-being (NEHW) study. *Health and Place*, 31, 65–74. <http://dx.doi.org/10.1016/j.healthplace.2014.11.001>
- O'Campo, P., Xue, X., Wang, M. C., & Caughy, M. (1997). Neighborhood risk factors for low birthweights in Baltimore: A multilevel analysis. *American Journal of Public Health*, 87, 113–118.
- O'Neil, R., Parke, R. D., & McDowell, D. J. (2001). Objective and subjective features of children's neighborhoods: Relations to parental regulatory strategies and children's social competence. *Applied Developmental Psychology*, 22, 135–155. [http://dx.doi.org/10.1016/S0193-3973\(01\)00073-9](http://dx.doi.org/10.1016/S0193-3973(01)00073-9)
- Ocaña-Riola, R., Saurina, C., Fernandez-Ajuria, A., Sánchez-Cantalejo, C., Saez, M., Ruiz-Ramos, & Benach, J. (2008). Area deprivation and mortality in the provincial capital cities of Andalusia and Catalonia (Spain). *Journal of Epidemiology & Community Health*, 62, 147–152. <http://dx.doi.org/10.1136/jech.2006.053280>
- Ogders, C. L., Caspi, A., Bates, C. J., Sampson, R. J., & Moffitt, T. E. (2012). Systematic social observation of children's neighborhoods using google street view: A reliable and cost-effective method. *Journal of Child Psychology and Psychiatry*, 53, 1009–1017. <http://dx.doi.org/10.1111/j.1469-7610.2012.02565.x>

- Park, R., Burgess, E. W., & McKenzie, R. D. (1925). *The city*. Chicago: University of Chicago Press.
- Pearl, M., Braveman, P., & Abrams, B. (2001). The relationship of neighborhood socioeconomic characteristics to birthweight among 5 ethnic groups in California. *American Journal of Public Health, 91*, 1808–1814. <http://dx.doi.org/10.2105/AJPH.91.11.1808>
- Perkins, D. D., Meeks, J. W., & Taylor, R. B. (1992). The physical environment of street blocks and resident perceptions of crime and disorder: Implications for theory and measurement. *Journal of Environmental Psychology, 12*, 21–34. [http://dx.doi.org/10.1016/S0272-4944\(05\)80294-4](http://dx.doi.org/10.1016/S0272-4944(05)80294-4)
- Perkins, D. D., & Taylor, R. B. (1996). Ecological assessments of community disorder: Their relationship to fear of crime and theoretical implications. *American Journal of Community Psychology, 24*, 63–107. <http://dx.doi.org/10.1007/BF02511883>
- Pinchevsky, G. M., & Wright, E. M. (2012). The impact of neighborhoods on intimate partner violence and victimization. *Trauma Violence Abuse, 13*, 112–132. <http://dx.doi.org/10.1177/1524838012445641>
- Quick, M. (2013). *Exploring crime in Toronto, Ontario with applications for law enforcement planning: Geographic analysis of hot spots and risk factors for expressive and acquisitive crimes* (unpublished master's thesis for master's degree). University of Waterloo, Ontario, Canada.
- Raudenbush, S., & Sampson, R. J. (1999). 'Ecometrics': Toward a science of assessing ecological settings, with application to the systematic social observation of neighborhoods. *Sociological Methodology, 29*, 1–41.
- Raykov, T. (2004). Behavioral scale reliability and measurement invariance evaluation using latent variable modeling. *Behavioral Therapy, 35*, 299–331. [http://dx.doi.org/10.1016/S0005-7894\(04\)80041-8](http://dx.doi.org/10.1016/S0005-7894(04)80041-8)
- Reiss, A. J. (1971). *The police and the public*. New Haven, CT: Yale University Press.
- Robinson, J. B., Lawton, B. A., Taylor, R. B., & Perkins, D. D. (2003). Multilevel longitudinal impacts of incivilities: Fear of crime, expected safety, and block satisfaction. *Journal of Quantitative Criminology, 19*, 237–274. doi: 10.1023/A:1024956925170.
- Roosa, M. W., Deng, S., Ryu, E., Burrell, G. L., Tein, J. Y., Jones, S., & Crowder, S. (2005). Family and child characteristics linking neighborhood context and child externalizing behavior. *Journal of Marriage and Family, 67*, 515–529. <http://dx.doi.org/10.1111/j.0022-2445.2005.00132.x>
- Ross, C. E., & Mirowsky, J. (1999). Disorder and decay: The concept and measurement of perceived neighborhood disorder. *Urban Affairs Review, 34*, 412–432. <http://dx.doi.org/10.1177/107808749903400304>
- Ross, C. E., & Mirowsky, J. (2001). Neighborhood disadvantage, disorder and health. *Journal of Health and Social Behavior, 42*, 258–276. <http://dx.doi.org/10.2307/3090214>
- Ross, C. E., & Mirowsky, J. (2009). Neighborhood disorder, subjective alienation, and distress. *Journal of Health and Social Behavior, 50*, 49–64. <http://dx.doi.org/10.1177/002214650905000104>
- Ross, C. E., Mirowsky, J., & Pribesh, S. (2001). Powerlessness and the amplification of threat: Neighborhood disadvantage, disorder, and mistrust. *American Sociological Review, 66*, 568–591.
- Sampson, R. J. (2009). Disparity and diversity in the contemporary city: Social (dis)order revisited. *The British Journal of Sociology, 60*, 1–31. <http://dx.doi.org/10.1111/j.1468-4446.2009.01211.x>
- Sampson, R. J. (2012). *Great American city: Chicago and the enduring neighborhood effect*. Chicago, IL: University of Chicago Press.
- Sampson, R. J., & Raudenbush, S. W. (1999). Systematic social observation of public spaces: A new look at disorder in urban neighborhoods. *American Journal of Sociology, 105*, 603–651. <http://dx.doi.org/10.1086/210356>
- Sampson, R. J., & Raudenbush, S. W. (2004). Seeing disorder: Neighborhood stigma and the social construction of broken windows. *Social Psychology Quarterly, 67*, 319–342. <http://dx.doi.org/10.1177/019027250406700401>
- Sampson, R. J., Raudenbush, S. W., & Earls, F. (1997). Neighborhoods and violent crime: A multilevel study of collective efficacy. *Science, 277*, 918–924. <http://dx.doi.org/10.1126/science.277.5328.918>
- Schaefer-McDaniel, N., Caughy, M. O., O'Campo, P., & Gearey, W. (2010). Examining methodological details of neighbourhood observations and the relationship to health: A literature review. *Social Science & Medicine, 70*, 277–292. <http://dx.doi.org/10.1016/j.socscimed.2009.10.018>
- Shaw, C. R., & McKay, H. D. (1942). *Juvenile delinquency and urban areas*. Chicago: University of Chicago Press.
- Skogan, W. G. (1986). Fear of crime and neighbourhood change in communities and crime. *Crime and Justice: A Review of Research, 8*, 203–229.
- Skogan, W. G. (1990). *Disorder and decline: crime and the spiral of decay in American cities*. Berkeley: University of California Press.
- Skogan, W. G., & Maxfield, M. (1981). *Coping with crime: Individual and neighborhood reactions*. Beverly Hills, Calif: Sage Publications.
- Sparks, C. S. (2011). Violent crime in San Antonio, Texas: An application of spatial epidemiological methods. *Spatial and Spatio-temporal Epidemiology, 2*, 301–309. <http://dx.doi.org/10.1016/j.sste.2011.10.001>
- Summers, A. A., Cheshire, P. C., & Senn, L. (1999). *Urban change in the United States and western Europe: Comparative analysis and policy*. Washington, D.C: The Urban Institute Press.
- Taylor, R. B. (1997). Social order and disorder of street blocks and neighborhoods: Ecology, microecology, and the systemic model of social disorganization. *Journal of Research in Crime and Delinquency, 34*, 113–155. <http://dx.doi.org/10.1177/0022427897034001006>
- Taylor, R. B. (2001). *Breaking away from broken windows: Evidence from Baltimore neighborhoods and the nationwide fight against crime, grime, fear and decline*. New York, NY: Westvie.
- Taylor, R. B. (2005). The Incivilities or "broken windows" thesis. In L. E. Sullivan (Ed.), *Encyclopedia of Law Enforcement*. Thousand Oaks: Sage.
- Taylor, R. B., & Shumaker, S. A. (1990). Local crime as a natural hazard: Implications for understanding the relationship between disorder and fear of crime. *American Journal of Community Psychology, 18*, 619–642. <http://dx.doi.org/10.1007/BF00931234>
- Tendulkar, S. A., Buka, S., Dunn, E. C., Subramanian, S. V., & Koenen, K. C. (2010). A multilevel investigation of neighborhood effects on parental warmth. *Journal of Community Psychology, 38*, 557–573. <http://dx.doi.org/10.1002/jcop.20381>
- Toet, A., & van Schaik, M. G. (2012). Effects of signals of disorder on fear of crime in real and virtual environments. *Journal of Environmental Psychology, 32*, 260–276. <http://dx.doi.org/10.1016/j.jenvp.2012.04.001>
- Veysey, B. M., & Messner, S. F. (1999). Further testing of social disorganization theory: An elaboration of Sampson and Groves's "community structure and crime". *Journal of Research in Crime and Delinquency, 36*, 156–174. <http://dx.doi.org/10.1177/0022427899036002002>
- Wei, E., Hipwell, A., Pardini, D., Beyers, J. M., & Loeber, R. (2005). Block observations of neighbourhood physical disorder are associated with neighbourhood crime, firearm injuries and deaths, and teen births. *Journal of Epidemiology & Community Health, 59*, 904–908. <http://dx.doi.org/10.1136/jech.2004.027060>
- White, R. M. B., Roosa, M. W., Weaver, S., & Nair, R. L. (2009). Cultural and contextual influences on parenting in Mexican American families. *Journal of Marriage and Family, 71*, 61–79.
- Wilson, W. J. (1987). *The truly disadvantaged: The inner city, the underclass and public policy*. Chicago: University of Chicago Press.
- Wilson, J. Q., & Kelling, G. (1982). Broken windows: The police and neighborhood safety. *Atlantic Monthly, 249*, 29–38.
- Winkleby, M. A., & Cubbin, C. (2003). Influence of individual and neighbourhood socioeconomic status on mortality among black, Mexican-American, and white women and men in the United States. *Journal of Epidemiology and Community Health, 57*, 444–452. doi: 10.1136/jech.57.6.444.
- Worton, S. K., Caplan, R., Nelson, G., Pancer, S. M., Loomis, C., Peters, R. D., & Hayward, K. (2014). Better Beginnings, Better Futures: Theory, research, and knowledge transfer of a community-based initiative for children and families. *Psychosocial Intervention, 23*, 135–143. <http://dx.doi.org/10.1016/j.psi.2014.02.001>
- Wright, E. M., & Benson, M. L. (2011). Clarifying the effects of neighborhood disadvantage and collective efficacy on violence "behind closed doors". *Justice Quarterly, 28*, 775–798. <http://dx.doi.org/10.1080/07418825.2010.533687>
- Xu, Y., Fiedler, M. L., & Flaming, K. H. (2005). Discovering the impact of community policing: The broken windows thesis, collective efficacy, and citizens' judgement. *Journal of Research in Crime and Delinquency, 42*, 147–186. <http://dx.doi.org/10.1177/0022427804266544>
- York Cornwell, E., & Cagney, K. A. (2014). Assessment of neighborhood context in a nationally representative study. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences, 69*, S51–S63. <http://dx.doi.org/10.1093/geronb/gbu052>