



Anuario de Psicología Jurídica 2024

<https://journals.copmadrid.org/apj>



Is the Self-reported Information by Male Sentenced of Violence against Women in the Intervention Phase Valid?

Ramón Arce¹, Verónica Marcos¹, Jéssica Sanmarco¹, and Francisca Fariña²

¹University of Santiago de Compostela, Spain; ²University of Vigo, Spain

ARTICLE INFO

Article history:
Received 30 May 2022
Accepted 13 October 2022
Available online 30 March 2023

Keywords:
Malingering
Batters
Defensiveness
Forensic evaluation
Denial of symptoms

A B S T R A C T

The evaluation of the needs of convicts is a critical issue for the design of the intervention. Nonetheless, in the information reported by convicts in penitentiary evaluation, simulation (malingering) and dissimulation (denial of symptoms, social desirability, and both) must be suspected. Thus, a field study with the aim of estimating the prevalence rate of the distortions and its effects in the needs reported by convicts was designed. A total of 624 men convicted for intimate partner violence reported their needs in the MMPI-2 in the reception phase of a penitentiary intervention program. The results exhibited consistent and non-malingered answers. Nevertheless, a significant prevalence of dissimulation consisting in denial of symptoms, .112 [.087, .137], social desirability, .223 [.190, .256], and the combination of both two, .120 [.095, .145]. The effects in denial of symptoms emerged in the information of less depressive (i.e., lack of interest in activities, excessive sensitivity, lack of sociability), paranoid (i.e., interpersonal sensitivity and perceptiveness), psychasthenic (i.e., general subjective distress and negative affect), schizoid (i.e., alienation, chaotic interpersonal relations, and disruptive cognitive processes), and hypomanic (i.e., hyperactivity, irritability, and sensation seeking) symptoms. Likewise, social desirability dissimulators detailed significantly less depressive, psychopathic (i.e., less general social maladjustment), paranoid, psychasthenic, schizoid, and hypomanic symptoms. Similarly, the whole dissimulators (combining denial of symptoms and social desirability responses) informed less depressive, psychopathic, paranoid, psychasthenic, schizoid, and hypomanic symptoms, and fewer problems of social introversion. Conversely, they related significantly more symptoms of hysteria (i.e., physical complaints). Distortions introduced by convicts in self-reports must be controlled to design effective correctional interventions.

¿Es válida la información reportada por sentenciados por maltrato contra la mujer en la fase de intervención?

R E S U M E N

La evaluación de las necesidades de los sentenciados es clave para diseñar la intervención. No obstante, en la información facilitada por los sentenciados en proceso de evaluación penitenciaria ha de sospecharse simulación y disimulación (negación de síntomas, deseabilidad o ambas). Por ello, se diseñó un estudio de campo con el objetivo de estimar la tasa de prevalencia de distorsiones en las respuestas y sus efectos en las necesidades psicológicas informadas. Una muestra de 624 hombres sentenciados por violencia de género informaron de sus necesidades en el MMPI-2 en la fase de acogida de un programa de intervención penitenciario. Los resultados pusieron de manifiesto que las respuestas de los sentenciados eran coherentes y no estaban sujetas a simulación. No obstante, se observó una prevalencia significativa de disimulación consistente en negación de síntomas, .112 [.087, .137], deseabilidad social, .223 [.190, .256], y la combinación de ambas, .120 [.095, .145]. Los efectos de las distorsiones en la negación de síntomas se manifestaron en el informe de menos síntomas depresivos (i.e., ocultación de carencias en la sociabilidad), paranoides (i.e., ocultación de sensibilidad interpersonal), psicasténicos (i.e., ocultación de afecto negativo), esquizoides (i.e., ocultación de procesos cognitivos disruptivos) e hipomaniacos (i.e., ocultación de irritabilidad y búsqueda de sensaciones). Asimismo, los efectos de la deseabilidad se concretaron en el informe de menos de síntomas de depresión, psicopatía (i.e., menos inadaptación social), paranoia, psicastenia, esquizofrenia e hipomanía. Igualmente, los disimuladores que combinaron deseabilidad social y negación de síntomas manifestaron menos rasgos depresivos, psicopáticos, paranoides,

Palabras clave:
Simulación
Maltratadores
Estilo de respuesta defensivo
Evaluación forense
Negación de síntomas

Cite this article as: Arce, R., Marcos, V., Sanmarco, J., & Fariña, F. (2024). Is the self-reported information by male sentenced of violence against women in the intervention phase valid? *Anuario de Psicología Jurídica*, 34, 23-30. <https://doi.org/10.5093/apj2023a3>

Funding: This research has been partially sponsored by a grant of the Consellería de Cultura, Educación e Ordenación Universitaria of the Xunta de Galicia (ED431B 2020/46), and by a grant of the Ministry of Science and Innovation of Spain (PID2020-115881RB-I00). Correspondence: ramon.arce@usc.es (R. Arce).

ISSN: 1133-0740/© 2024 Colegio Oficial de la Psicología de Madrid. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

psicasténicos e hipomaniacos, así como menos problemas de introversión social. Además, notificaron más histeria (i.e., quejas somáticas). Las distorsiones introducidas por los convictos en los auto-informes deben ser controladas para el diseño de intervenciones penitenciarias eficaces.

In psychological evaluation, the clinician must suspect simulation¹, understood as “the intentional production of false or grossly exaggerated physical or psychological symptoms, motivated by external incentives such as avoiding military duty, avoiding work, obtaining financial compensation, evading criminal prosecution, or obtaining drugs” (American Psychiatric Association [APA, 2013, p. 726]), when a combination of the following is noted: a) evaluations in medicolegal context; b) a marked discrepancy between the alteration manifested by the person and the objective data obtained from the clinical examination; c) lack of cooperation during the diagnostic evaluation and non-compliance with the prescribed treatment regimen; and d) the presence of antisocial personality disorder. Nevertheless, in the forensic setting, simulation must be suspected in any circumstance in which the evaluated person can obtain a benefit from presenting a psychological disorder, forcing a simulation differential diagnosis, such that if this cannot be ruled out the psychological evaluation is not valid as forensic evidence of charge (Arce, 2018) or in criminal insanity evaluations (Osuna et al., 2015).

Additionally, in the forensic psychological evaluation not only should simulation be suspect, but, as the case may be, dissimulation² and oversimulation³ (exaggeration of symptoms). Oversimulation is the exaggeration of a real psychological damage to obtain a benefit, generally economic, as in the case of compensation for the aftermath of a traffic accident. Meanwhile, dissimulation consists of reporting responses biased towards defensiveness. This includes the denial or concealment of symptoms – negative dissimulation – and the social desirability – positive dissimulation –, that is, the assumption of positive characteristics that are not present (fabricated) or the concealment of negative characteristics to convey a favorable image of oneself, being able to be conscious – impression management – or unconscious – self-deception – (Paulhus, 1984). Dissimulation must be suspected in the evaluation of populations in intervention for behavior problems or for violent behaviors (Molero et al., 2022; Nasaescu et al., 2020); of prison inmates (Arce et al., 2014; Martín et al., 2019; Redondo et al., 2022); in risk assessment (Jugl et al., 2021); and in the forensic evaluation of family courts (Arce et al., 2015; Baer & Miller, 2002). Overall, dissimulation is more difficult to classify correctly than simulation (Baer & Miller, 2002), because the task involves less cognitive activity of the individual and, consequently, less measure of the construct (validity) (Arce et al., 2015). These response biases, if they are systematic, constitute a serious measurement problem, since they support an interpretive hypothesis of the relationship between the variables that is different from the real one (Podsakoff et al., 2003).

In sum, forensic evaluation requires a psychological diagnosis (variable according to judicial needs) and, at the same time, a differential diagnosis of distortion of responses (simulation, dissimulation and/or oversimulation). With this dual purpose, the traditional clinical evaluation is not valid, since it does not diagnose simulation (the diagnosis or suspicion of dissimulation and oversimulation do not exist in clinical setting; APA, 2013), due to the potentially devastating consequences of a false positive: deny a treatment to a sick person (it could lead to judicial responsibility), which may be a plea for help or of a phenomenological style to overreact and to be traumatized (Greene, 2011; Rogers, 2018). On the contrary, in the forensic psychological evaluation the undesirable outcome would be a false negative of simulation, dissimulation, or oversimulation. For the study and classification of response biases, both individual and population, a multi-method approach is required (Burchett & Bagby, 2014; Graham, 2011) comprising an interview,

necessary for the forensic diagnosis and the establishment of causality (Arce, 2017), and a psychometric measure, with which to contrast –inter- measure consistency – the forensic diagnosis and the differential diagnosis of distortion of responses (Graham, 2011; Greene, 2011).

The clinical-forensic interviews in use, i.e., the Structured Inventory of Reported Symptoms (SIRS; Rogers et al., 1992) and the Forensic-Clinical Interview (Vilarinho et al., 2013), embrace measures of the distortion of the responses (i.e., detection strategies for simulation, dissimulation and oversimulation), but do not have sufficient validity (i.e., false positive diagnosis is no 0) for the differential diagnosis of simulation, dissimulation, or oversimulation in the forensic context. With this purpose, the psychometric evaluation is fundamental, being the MMPI-2 the reference instrument in forensic setting (Graham, 2011; Greene, 2011) and with superiority over its Restructured Form, the MMPI-2-RF (Fariña et al., 2017; Gancedo et al., 2020; Redondo et al., 2019; Rogers et al., 2003; Sharf et al., 2017). In this regard, the MMPI-2 has numerous scales (in the forensic context, not only the scales of the commercial version are taken, but also other additional ones) and indices that have been extensively studied, showing significant discriminative efficacy and a great magnitude in the classification of simulation and oversimulation (Gancedo et al., 2020; Rogers et al., 2003), as well as dissimulation (Baer & Miller, 2002; Fariña et al., 2017).

The prodigality of scales and indices is necessary for two reasons. First, no scale or index by itself correctly classify the distortions of the responses, so the combination of scales and indices increase the validity of the approach to the control of false positives (forensic task; Fariña et al., 2014). Second, (over)simulation and dissimulation are not a unitary construct, but rather hide different strategies. Thus, (over)simulation can manifest itself in rare symptoms (*F_p* scale), quasi-rare (*F* and *F_b* scales), combination of symptoms (the family of *F* scales), severity of symptoms, or exaggeration of symptoms, i.e., overstimulation (the family of *F* scales). Hence, measures of all these strategies must be taken, combined in a measure of (over)simulation: infrequency (Gancedo, Sanmarco et al., 2021; Greene, 2011; Rogers, 2018; Vilarinho et al., 2013). This (over)simulation measure is completed by the lack of consistency in the responses (APA, 2013), which are evaluated by the *TRIN* and *VRIN* scales, as well as the *F-F_b* index (Fariña et al., 2014; Greene, 2011). Conversely, dissimulation can manifest itself in denial or concealment of symptoms, which are evaluate by the *K*, *L* and *S* scales and the Gough index (Butcher & Han, 1995); and social desirability (only referring to the conscious impression management, since non-conscious self-deception is part of the personality of the evaluated person) measure by the *L*, *MP*, *Od*, and *Wsd* scales (Arce et al., 2015; Bagby & Marshall, 2004; Strong et al., 1999).

Penitentiary intervention is aimed at addressing the criminogenic needs of inmates (i.e., inmate characteristics that contribute to the likelihood of recidivating) (Bonta & Andrews, 2017), such that if it is not adequately adjusted to the needs of each individual, it is not effective (Bonilla-Algobia & Rivas-Rivero, 2022; Moral-Jiménez & González-Sáez, 2020; Santirso et al., 2020) and can even be counterproductive (Arias et al., 2020). Hence, inmate needs are evaluated in the reception phase of the intervention with the aim of establishing the contents of the intervention programs based on the evaluation and adjusted to each case. This assessment relies almost exclusively on self-reports of moot reliability to prescribe treatment based on them. Thus, they are not consistent with other more reliable measures such as judicial or prison records (Expósito-Álvarez et al., 2021).

In this state of the question, a field study with a sentenced population was designed to find out the rate of prevalence of distortions in the responses and their effects, where appropriate, on the reported psychological needs (Arce et al., 2014).

Method

Participants

A total of 624 men between the ages of 18 and 78 ($M = 41.87$, $SD = 12.12$) serving a sentence of less than two years in prison, suspended by participation in a community penitentiary intervention program, participated in the study. All of them were gender-based primary offenders whose sentence was suspended by a voluntary participation in a community penitentiary program in application of L.O. 1/2004 (Ley Orgánica 1/2004, de 28 de diciembre, de Medidas de Protección Integral contra la Violencia de Género). The sentence suspension time was 5 years for 13 (2.1%), 4 years for 24 (3.8%), 3 years for 210 (33.7%), and 2 years for 377 (60.4%) of the batterers.

Procedure and Design

The sentenced persons were offered, under the protection of article 83 of L.O. 1/2015 (Ley Orgánica 1/2015 de 30 de marzo, por la que se modifica la Ley Orgánica 10/1995, de 23 de noviembre, del Código Penal), the suspension of the execution of the sentence in prison for participation in the Galician Re-Education Programme for Male Intimate-Partner Violence Offenders (Arce & Fariña, 2006, 2010). All participated voluntarily and signed the informed consent for the evaluation and intervention. In the reception phase, and after presenting the program and signed the acceptance form of participation, they were evaluated on the psychological adjustment. The order in which measures (i.e., assessment of psychological adjustment, risk assessment, assessment of cognitive skills, assessment of psychosocial skills) were administered to participants, was rotated, following a standard systematic rotation procedure (i.e., A, B, C, D,; B, C, D,; C,), in order to control the carry-over effect of the measurement variables in the subsequent ones (Arce et al., 2000).

Measurement Variables

Participants were administered the Spanish adaptation of the MMPI-2 (Butcher et al., 2019), the reference instrument for evaluation in forensic and penitentiary settings (Ackerman & Pritzl, 2011; Arce et al., 2015; Archer et al., 2006; Fariña et al., 2017; Graham, 2011; Greene, 2011; Redondo et al., 2019; Rogers et al., 2003). This instrument measures the psychological adjustment of the individual in 10 basic clinical scales: Hypochondriasis [Hs], Depression [D], Hysteria [Hy], Psychopathic Deviate [Pd], Paranoia [Pa], Psychasthenia [Pt], Schizophrenia [Sc], Hypomania [Ma], and Social Introversion [Si]. The Masculinity-Femininity scale was discarded since it did not measure psychological adjustment (Graham, 2011). This questionnaire includes also scales for the differential diagnosis of distortions in responses. With the measures and validity scales, 5 response manipulation profiles were constructed in which, due to the addition of various measures, the probability of a distortion of the responses is very high: inconsistent responses, infrequent responses (over-simulation), denial of symptoms (negative dissimulation), social desirability (positive dissimulation), and dissimulation (whole dissimulation: negative and positive dissimulation).

The APA (2013) include as a criterion for the suspicion of simulation the lack of consistency in the responses (that is, in non-psychiatric populations it evaluates simulation). This is measured in the MMPI-2 (Greene, 2011) through the VRIN (variable response inconsistency)

and TRIN (true response inconsistency) scales to which the |F-Fb| index has been added, which assesses the infrequency of symptoms in the first part of the instrument (up to item 370; included the basic clinical scales), and Fb which is the continuation of F in the second part (from item 370 onwards). As a criterion for the suspicion of a very high probability of inconsistency in the responses, two of these indicators ($VRIN \geq 9$, $TRIN \geq 11$; $|F-Fb| > 11$) have to classify the protocol as 'inconsistent' (Osuna et al., 2015). Simulation, which is measured through the infrequency of symptoms (Gancedo, Sanmarco et al., 2021; Greene, 2011; Rogers, 2018), that is, symptoms that rarely occurred in the general population, is evaluated with the scales F (Infrequency; items-symptoms-that are infrequently admitted by the normative population, < 10%), Fb (Back Infrequency scale; continuation of F on the back of the instrument); Fp (Uncommon psychopathology; items-symptoms-that are infrequently admitted by psychiatric inmates, < 20%), and the Gough index (i.e., F-K), being highly probable that the responses are distorted when two or more of the following criteria are met: $F, Fb, Fp \geq T66.45$; and Gough index ≥ 30 (Arce et al., 2009; Rogers, 2018).

Denial or concealment of symptoms (favorable report of psychopathology) is addressed with K (correction scale; people who create a favorable impression of themselves; denial of symptoms especially in somatization and histrionic disorders), L (lie; intent deliberate not to respond honestly and frankly, including denial of minor misconduct, honesty, aggressiveness, bad thoughts, and weakness of character), S (superlative; evaluates people who are superlative) scales and the Gough index (Butcher & Han, 1995). A protocol is classified as highly probable subject to denial of symptoms when two or more of the following criteria are met: $K < 40$, $L \geq 66.45$, $S \geq 66.45$, Gough index < -21 (Greene, 2011; Fariña et al., 2017). The social desirability in the responses (that was, the intentional handling of the responses in order to make a positive impression) is appreciated with the scales L, Mp (positive malinger; identifies self-favorable protocols of psychopathology), Od (other deception; update of Mp) and Wsd (Wigging's social desirability scale; responses biased to appear socially desirable), and the classification as very highly probable impression management when two or more of the criteria are met: $L, MP, Od, Wsd \geq T66.45$ (Arce et al., 2015).

Data Analysis

For the study of the prevalence rate of response distortion strategies, the observed probability was contrasted with a constant, a trivial probability (insignificant, .05; Fandiño et al., 2021), calculating the zeta score for the difference of the observed proportion with a constant. β was computed in order to know the power of the results, with $1-\beta$ being the probability associated with the correct refutation of the null hypothesis. The effect size was measured in odds ratio (OR) and the magnitude was quantified in terms of the effect incremental index (EII; Redondo et al., 2019) that estimates the increased probability over the constant due to the distortion strategy. As to compare the observed probabilities in the different strategies, 95% confidence intervals for the probabilities were computed: if confidence intervals overlap, the observed probabilities are equal, and if the confidence intervals do not overlap, the observed probabilities are significantly different ($p < .05$).

A comparison of means with MANOVAs with the psychological adjustment measurement variables as dependent variables ($\bar{r} = .331$, $p < .001$) and the group (distorted response pattern vs. undistorted response pattern) as a factor was applied. The analysis of variance is robust to the heterogeneity of variances when the groups are of equal size (i.e., $N_{\text{greater}}/N_{\text{lesser}} < 1.5$), but when the assumption of homogeneity of variances is violated and the Ns are different, it could result in relevant deviations in the estimate of the associated probability (α) to F (Stevens, 1986). In multivariate F-test, the Pillai-

Bartlett (V) trace is robust to the heterogeneity of variance (Olson, 1976), but in univariate *F*-tests, when the variability is greater in the larger group, *F* is conservative and vice versa. To address this contingency and have absolute certainty of the correct rejection of the null hypothesis, three safeguards were verified (Mayorga et al., 2020): a) that the theoretical *F* (Box's conservative test, 3.841) was less than the empirical value of *F*; b) that the type II error – false acceptance of the null hypothesis – was greater than the type I error – false rejection of the null hypothesis – i.e., $\beta/\alpha > 1$, being the result the number of times that the probability of a false negative than a false positive; and c) that the effect size, corrected for unequal *N*s – the analysis of variance is robust to the heterogeneity of variance with equal *N*s –, was > 0.20 (small). In all results with significant univariate *F*-tests ($p < .05$), these 3 requirements were met.

The effect size for multivariate comparisons was estimated with η^2_p , interpreted in terms of the explained variance and quantified as the probability of superiority of the effect size (PS_{ES} ; Gancedo, Fariña et al., 2021). For the difference of two means, a standardized mean difference, *d*, was calculated with the Hedges' formula (when $N1 \neq N2$), interpreting it in terms of the PS_{ES} . From *d*, the quantification of the increase (+) or decrease (-) in the distortions in the responses was estimated with an adaptation of the BESD (Rosenthal & Rubin, 1982), *r* (Gancedo, Fariña, et al., 2021). Finally, the error of the statistical model through the probability of an inferiority score (PIS; Fandiño et al., 2021) was calculated, which is an estimate of the probability that an individual in the group, in our case of distorted responses, obtained a lower score in the measurement variable than the mean of the group of non-distorted responses (that is, probability contrary to the classification of the statistical model).

Ethical Issues

The data were taken anonymized from the database of the Galician Re-Education Programme for Male Intimate-Partner Violence Offenders and were treated guaranteeing all the rights of convicted set forth in Penitentiary Spanish Law (Ley Orgánica 1/1979, de 26 de septiembre, General Penitenciaria) and according with the Spanish Data Protection Law (Ley Orgánica 3/2018, de 5 de diciembre, de Protección de Datos Personales y Garantía de los Derechos Digitales).

Results

Study of the Validity of the Protocols

The protocols were scrutinized to identify whether they were subject to extreme acquiescence ($TRIN T \geq 80$), a pattern of random responses ($VRINT \geq 80$; F, Fb or $Fp T \geq 120$), or outliers ($Lrs > 12$ or $Krs > 29$ i.e., 99.9 percentile), to eliminate them from the database (Fariña et al., 2014; Graham, 2011; Greene, 2011). Lack of collaboration (> 10

unanswered items or with double response; Graham, 2011) was also stated as a cause of invalidity of the protocols in forensic or prison evaluation. No protocol was removed for these reasons.

Prevalence of Cases of Distorted Responses

The results (see Table 1) showed a significantly negative rate (lower than normative, .05) of inconsistent responses among those convicted of intimate-partner violence, indicating that the responses of male convicted of intimate-partner violence were more consistent than the normative. Likewise, the response profiles of offenders were not uncommon, because the responses of these individuals do not include symptoms that rarely occur in the general ($< 10\%$) or psychiatric ($< 20\%$) population. In sum, in this evaluation context this population did not simulate or exaggerate psychopathology. On the contrary, the observed rates of denial of symptoms (negative dissimulation), social desirability (positive dissimulation), and whole dissimulation (combination or concealment of denial of symptoms and social desirability) have been registered significantly in this population, suggesting that the observed prevalence was not trivial. The magnitude of the effect of the distortions in the responses introduced by the subjects was moderate ($OR \approx 2.47$) in whole dissimulation and denial of symptoms, and large ($OR > 4.25$) in social desirability, being the increase in the magnitude of the effect (EII) due to the conscious distortions in the responses of 55.4% in denial of symptoms, 58.3% in dissimulation, and 77.6% in social desirability. Comparatively, the prevalence of distorted responses in the direction of social desirability was significantly higher (the lower limit of the confidence interval for the observed prevalence is higher than the upper limit of denial of symptoms and dissimulation) than that of denial of symptoms and whole dissimulation.

As a consequence of these results, it was necessary to contrast what clinical symptoms they deny or conceal, and what positive characteristics report they did not present (social desirability), as well as to quantify the magnitude of the response distortions.

Negative Dissimulation: Denial of Symptoms

Knowing that prevalence rate of denial of symptoms was significant, the results showed that denial has significant effects on the self-reports of psychological adjustment, $F(9, 605) = 8.39, p < .001, 1-\beta = 1.0$, accounting for the 11.1% of the variance ($\eta^2_p = .111$), an effect size greater than 69.15% of all effect sizes ($PS_{ES} = .6915$).

The univariate effects (see Table 2) exhibited that batterers classified as 'denial of symptoms' reported significantly less depressive (i.e., lack of interest in activities expressed as general apathy, physical symptoms such as sleep disturbances and gastrointestinal ailments, excessive sensitivity, and lack of sociability), paranoid (i.e., interpersonal sensitivity and perceptiveness), psychasthenic (i.e., general subjective distress and negative affect), schizoid – in non-psychiatric populations the Schizophrenia scale measures social alienation, not the positive symptoms of schizophrenia – (i.e.,

Table 1. Prevalence of Distorted Responses

Distorted responses	$f(p)$ [95% CI_p]	Z	1- β	OR	EII ¹
Inconsistency	8(.013) [.004, .022]	-4.24***	.999	0.26	-.740
Infrequency(simulation)	41(.066) [.047, .085]	1.83	.535	1.32	.242
Denial of symptoms	69(.112) [.087, .137]	7.05***	.999	2.44	.554
Social desirability	139(.223) [.190, .256]	19.80***	1.00	4.46	.776
Dissimulation	75(.120) [.095, .145]	8.02***	.999	2.40	.583

Note. $f(p)$ [95% CI_p] = frequency(proportion) [95% confidence interval]; Z = zeta score for the comparison of the observed probability in the batterer population with a constant (.05; trivial effect); OR = odds ratio; ¹EII = effect incremental index.

*** $p < .001$.

Table 2. Univariate Effects on the MMPI-2 Basic Clinical Scales for the Denial of Symptoms Desirability Factor. Between-subjects Effects

Clinical scales	<i>F</i>	<i>d</i>	1- β	M_{N-NS}	M_{NS}	PS_{ES}	PIS
Hs	0.60	0.10	.120	51.55	52.61	.055	.460
D	4.39*	-0.27	.553	53.71	50.99	.151	.394
Hy	6.95**	0.34	.749	50.89	54.67	.190	.367
Pd	1.35	-0.15	.213	54.58	53.09	.088	.440
Pa	26.12***	-0.65	.999	56.74	49.10	.354	.258
Pt	8.87**	-0.38	.844	50.00	46.32	.213	.352
Sc	7.13**	-0.34	.760	50.15	46.58	.190	.367
Ma	25.64***	-0.65	.999	51.97	45.70	.354	.258
Si	14.42***	-0.48	.966	50.87	46.58	.266	.316

Note. *df*(1, 613); M_{N-NS} = mean of the 'not-denial of symptoms group'; M_{NS} = mean of the 'denial of symptoms group'.
* $p < .05$, ** $p < .01$, *** $p < .001$.

alienation, chaotic interpersonal relations, and disruptive cognitive processes), and hypomanic (i.e., hyperactivity, irritability, and sensation seeking) symptoms. They also warned of less social introversion (less personal discomfort in social situations, isolation, general maladjustment, and self-deprecation) than those in the non-denial of symptoms group. Finally, and contrary to the grouping criterion, the batterers in the denial of symptoms group reported significantly more symptoms of hysteria (i.e., complaints about muscular, gastrointestinal or cardiac problems, feelings of tension, anguish or fear). Nevertheless, this measure includes many items that imply a denial of problems in life or the absence of social anxiety (Greene, 2011) that, indeed, are related to denial of symptoms. Thus, the measure is reversed, and the results corroborated the validity of the inmate classification model.

Quantitatively, the batterers of the denial of symptoms group reported, on average, a 13.4% ($r = -.134$) fewer depressive symptoms; 30.9% ($r = -.309$) fewer paranoid symptoms; 18.7% ($r = -.187$) fewer symptoms of psychasthenia; 16.8% ($r = -.168$) fewer schizoid symptoms; 30.9% ($r = -.309$) fewer hypomanic symptoms; and 23.3% ($r = -.233$) fewer problems related to social introversion. Furthermore, the batterers of the denial of symptoms group assumed 16.8% ($r = .168$) more characteristics of hysteria (denial of problems in life or the absence of social anxiety). The magnitude of the effect (PS_{ES}) was greater (damage in mental health markers) than 15.1% of all possible in depression; than 19.0% in hysteria; than 35.4% in paranoia; than 21.3% in psychasthenia; than 19.0% in schizophrenia; than 35.4% in hypomania; and than 26.6% in social introversion. On the contrary, the probability (PIS) that the batterers of the denial of symptoms group obtained a score lower than the mean of the non-denial of symptoms group (error of the statistical model) was 39.4% in depression; 25.8% in paranoia; 35.2% in psychasthenia; 36.7% in schizophrenia; 25.8% in hypomania; and 31.6% in social introversion; while in hysteria,

the probability that those in the non-denial of symptoms group had a lower score than those in the denial of symptoms group was of 36.7%.

Positive Dissimulation: Social Desirability

Based on previous findings supporting that the prevalence rate of social desirability in batterer responses was significant, the results displayed that the social desirability factor has significant effects on the self-reports of psychological adjustment, $F(9, 612) = 10.88$, $p < .001$, $1-\beta = 1.0$, accounting for 13.8 % of the variance ($\eta^2_p = .138$), an effect size greater than 71.57% of all effect sizes ($PS_{ES} = .7157$).

The univariate effects (see Table 3) revealed that batterers in the social desirability group reported significantly fewer depressive, psychopathic (i.e., general social maladjustment and absence of pleasant experiences), paranoid, psychasthenic, schizoid, and hypomanic symptoms.

Similarly, they reported less social introversion than those of the group of non-social desirability. Quantitatively, the batterers of the group who distorted their answers in line with social desirability reported, on average, 12.9% ($r = -.129$) fewer depressive symptoms; 25.6% ($r = -.256$) fewer psychopathic characteristics; 27.0% ($r = -.270$) fewer paranoid symptoms; 26.5% ($r = -.265$) fewer psychasthenic symptoms; 28.3% ($r = -.283$) fewer schizoid symptoms; 18.2% ($r = -.182$) fewer hypomanic symptoms; and 31.3% ($r = -.313$) fewer problems related to social introversion than batterers of the non-social desirability group. The magnitude of the effect was greater (damage in mental health markers) than 14.3% of all possible in depression; than 29.6% in psychopathic deviation; than 31.1% in paranoia; than 30.3% in psychasthenia; than 41.7% in schizophrenia; than 20.5% in hypomania; and than 36.2% in social introversion. On the contrary, the probability (PIS) that the batterers in the social desirability group obtained a score lower than the mean of the social undesirability group (statistical model error) was 39.7% in depression; 29.8% in

Table 3. Univariate Effects on the MMPI-2 Basic Clinical Scales for the Social Desirability factor. Between-subjects Effects

Clinical scales	<i>F</i>	<i>d</i>	1- β	M_{N-NS}	M_{NS}	PS_{ES}	PIS
Hs	2.74	-0.16	.380	51.92	50.21	.088	.436
D	7.46**	-0.26	.778	53.96	51.29	.143	.397
Hy	0.49	0.02	.056	51.14	51.38	.008	.492
Pd	33.08***	-0.53	1.00	55.59	50.14	.296	.298
Pa	34.23***	-0.56	1.00	57.30	50.71	.311	.288
Pt	32.36***	-0.55	1.00	50.72	45.49	.303	.291
Sc	37.78***	-0.59	1.00	51.08	45.01	.417	.278
Ma	15.11***	-0.37	.973	52.20	48.52	.205	.356
Si	47.44***	-0.66	1.00	51.63	45.93	.362	.255

Note. *df*(1, 613); M_{N-NS} = mean of the 'non-social desirability group'; M_{NS} = mean of the 'social desirability group'.
** $p < .01$, *** $p < .001$.

Table 4. Univariate Effects on the MMPI-2 Basic Clinical Scales for the Dissimulation Factor. Between-subjects Effects

Clinical scales	<i>F</i>	<i>d</i>	1- β	M_{N-D}	M_D	PS _{ES}	PIS
Hs	0.05	-0.03	.056	51.60	51.29	.016	.488
D	5.27*	-0.28	.630	53.70	50.83	.159	.390
Hy	5.99*	0.30	.686	50.81	54.19	.166	.382
Pd	7.97**	-0.35	.805	54.80	51.31	.197	.363
Pa	26.64***	-0.64	.999	56.73	49.27	.347	.261
Pt	11.13***	-0.41	.915	50.03	46.04	.228	.341
Sc	10.66***	-0.40	.903	50.24	46.03	.221	.345
Ma	28.89***	-0.66	1.00	52.13	45.69	.362	.255
Si	26.08***	-0.63	.999	50.99	45.49	.347	.264

Note. $df(1, 622)$; M_{N-D} = mean of group of 'non-dissimulators'; M_D = mean of group of 'whole dissimulators'.

* $p < .05$, ** $p < .01$, *** $p < .001$.

psychopathic deviation; 28.8% in paranoia; 29.1% in psychasthenia; 27.8% in schizophrenia; 35.6% in hypomania; and 25.5% in social introversion.

Whole Dissimulation

The results for the whole dissimulation factor (combination of denial or concealment of symptoms and social desirability) showed a significant effect on the self-reports of psychological adjustment, $F(9, 614) = 9.92$, $p < .001$, $1-\beta = 1.0$, accounting for 12.7% of the variance ($\eta^2_p = .127$), an effect size greater than 70.54% of all effect sizes ($PS_{ES} = .7054$).

The univariate effects (see Table 4) revealed that the batterers of the whole dissimulating group informed significantly fewer depressive, psychopathic, paranoid, psychasthenic, schizoid and hypomanic symptoms, and fewer problems of social introversion. In addition, they related significantly more symptoms of hysteria. Quantitatively, the batterers of the dissimulating group, on average, refer 14.0% ($r = -.140$) less of depressive symptoms; 17.2% ($r = -.172$); less of psychopathic characteristics; 30.5% ($r = -.305$) less of paranoid symptoms; 20.1% ($r = -.201$) less of symptoms of psychasthenia; 20.0% ($r = -.200$) less of schizoid symptoms; 31.3% ($r = -.313$) less of hypomanic symptoms; and 30.0% ($r = -.300$) less of problems related to social introversion. Additionally, the batterers of the group of dissimulating group assume 15.0% ($r = .150$) more characteristics of hysteria (denial of problems in life or the absence of social anxiety). The magnitude of the effect size was greater (i.e., damage in mental health markers) than 15.9% of all possible in depression; than 16.6% of all possible in hysteria; than 19.7% in psychopathic deviation; than 34.7% in paranoia; than 22.8% in psychasthenia; than 22.1% in schizophrenia; than 36.2% in hypomania; and than 34.7% in social introversion. On the contrary, the probability (PIS) that the batterers of the dissimulating group obtained a score higher than the mean of the non-dissimulating group (statistical model error) was of 39% in depression; of 36.3% in psychopathic deviation; 26.1% in paranoia; 34.1% in psychasthenia; 34.5% in schizophrenia; 25.5% in hypomania; and 26.4% in social introversion, while in hysteria the probability that those from the non-dissimulating group have a lower score under the mean of the dissimulators group was 38.2%.

Discussion

The results of this research are subjected to limitations in their generalization. First, the evaluation was carried out in the reception phase in the penitentiary treatment, so its generalization to other moments of the process is not automatic. Thus, while it was expected that during the penitentiary intervention the response biases would be in the direction of dissimulation, motivated by obtaining penitentiary benefits or therapeutic success, during the judicial

process a bias was expected in line with the simulation aimed at attenuation or exemption from criminal responsibility. Second, the results are not generalizable to other measurement instruments and subject classification criteria. Third, the magnitude of the effects cannot be generalized to other unmeasured criminogenic needs that may be the aim of penitentiary treatment.

Having these limitations in mind, the following conclusions emerged from the results. First, the conscious distortions in the responses of those evaluated in penitentiary intervention are consistent, suggesting that batterers manipulate the responses in a stable way throughout the measure. Hence, the lack of consistency in the responses must be taken as a characteristic of the individual and not as a measure of deception (Osuna et al., 2015). Second, in the evaluation in the intervention phase, (over)simulation should not be systematically suspected (i.e., rare, quasi-rare symptoms, severity of symptoms, combination of symptoms, oversimulation; Rogers, 2018; Vilarinho et al., 2013). Consequently, the information on deficiencies or needs must be taken, in general, as valid and, consequently, direct the intervention to them (Arce et al., 2014; Santirso et al., 2020). Third, in the evaluation of batterers in the intervention phase, dissimulation must be suspected, both referring to the denial or concealment of symptoms (non-criminogenic needs that are related to the criminal conduct but that are not the cause of it), as well as to social desirability (criminogenic needs, that are to say, causing the penalized conduct), or to a combination of both (whole dissimulation). Hence, the validity of the information self-reported by the batterers must be contrasted and complemented with other measures (Expósito-Álvarez et al., 2021). In sum, the validity of the measure is imperfect, understanding by imperfect validity that the results are not totally invalid, since not all subjects distort the responses (see the probability of cases contrary to the model; PIS). Fourth, manifested positive characteristics and unreported criminogenic needs (Bonta & Andrews, 2017) are those with the highest probability of bias being the self-report validity suspected. Failure to correct this bias will mean that the intervention does not focus on the needs of the individual and, henceforth, lost its effectiveness (Daffern et al., 2022). Fifth, there is also a systematic tendency to hide and deny clinical symptoms that, although they are not criminogenic needs (causal relationship with the criminal behavior), non-intervention on them is related to recidivism (Basanta et al., 2015). Sixth, the magnitude of the hidden or denied clinical symptoms, negative characteristics (e.g., denial of problems), and the invented positive characteristics range between approximately 1/5 and 1/3 of what is self-reported.

In conclusion, the effects of distortions introduced by batterers in self-reports must be taken into account in the design of both effective correctional interventions and future research with offenders.

Conflict of Interest

The authors of this article declare no conflict of interest.

Notes

¹This is the seminal scientific and judicial expression for intended fabrication of symptoms (Latin word with this significance) that was translated to English as malingering, feigning, feigning bad, faking, faking bad, overreporting, self-unfavorable reporting or even dissimulation. Some use them as synonymous, meanwhile others as a more specific definition of simulation (in any case, a simulation) and others as a non-intended (then it is not simulation). Thus, the correct and non-confusing expression for intended fabrication of symptoms is simulation.

²The seminal scientific and judicial expression (Latin word antonymous of simulation meaning the concealment of symptoms or negative characteristics including the assumption of non-present property to hide its lack) for defensiveness, feigning good, faking good, underreporting, self-favorable reporting.

³The seminal scientific and judicial expression (Latin word antonymous of simulation meaning the concealment of symptoms or negative characteristics including the assumption of non-present property to hide its lack) for defensiveness, feigning good, faking good, underreporting, self-favorable reporting.

References

- Ackerman, J. J., & Pritzl, T. B. (2011). Child custody evaluation practices: A 20-year follow-up. *Family Court Review*, 49(3), 618-628. <https://doi.org/10.1111/j.1744-1617.2011.01397.x>
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). American Psychiatric Association.
- Arce, R. (2017). Análisis de contenido de las declaraciones de testigos: evaluación de la validez científica y judicial de la hipótesis y la prueba forense [Content analysis of the witness statements: Evaluation of the scientific and judicial validity of the hypothesis and the forensic proof]. *Acción Psicológica*, 14(2), 171-190. <https://doi.org/10.5944/ap.14.2.21347>
- Arce, R. (2018). Evaluación del daño psicológico: psicometría, entrevista y técnica forense [Assessment of psychological harm: Psychometry, interview and forensic technique]. In E. Carbonell, D. Pineda, & M. Novo (Eds.), *Psicología jurídica: ciencia y profesión. Colección Psicología y Ley*, 15 (pp. 235-244). Sociedad Española de Psicología Jurídica y Forense. Retrieved from <http://sejpf.org/wp-content/uploads/2019/03/Vol.-15.-Psicolog%C3%ADa-Jur%C3%ADica.-Ciencia-y-profesi%C3%B3n.pdf>
- Arce, R., & Fariña, F. (2006). Programa Galicia de reeducación para maltratadores de género (Galician re-education programme for male intimate-partner violence offenders). *Anuario de Psicología Jurídica*, 16(1), 41-64. <https://journals.copmadrid.org/apj/archivos/102982.pdf>
- Arce, R., & Fariña, F. (2010). Diseño e implementación del Programa Galicia de reeducación de maltratadores: una respuesta psicosocial a una necesidad social y penitenciaria (Design and implementation of the Galician program for batterers re-education: A psychosocial answer to a social and penitentiary need). *Psychosocial Intervention*, 19(2), 153-166. <https://doi.org/10.5093/in2010v19n2a7>
- Arce, R., Fariña, F., Carballal, A., & Novo, M. (2009). Creación y validación de un protocolo de evaluación forense de las secuelas psicológicas de la violencia de género [Creation and validation of a forensic protocol to assess psychological harm in battered women]. *Psicothema*, 21(2), 241-247. Retrieved from <http://www.psicothema.com/pdf/3621.pdf>
- Arce, R., Fariña, F., & Fraga, A. (2000). Género y formación de juicios en un caso de violación [Gender and juror judgment making in a case of rape]. *Psicothema*, 12(4), 623-628. Retrieved from <http://www.psicothema.com/pdf/381.pdf>
- Arce, R., Fariña, F., & Novo, M. (2014). Competencia cognitiva en penados primarios y reincidentes: Implicaciones para la reeducación [Cognitive competence among recidivist and non-recidivist prisoners: Implications for the rehabilitation]. *Anales de Psicología*, 30(1), 259-266. <https://doi.org/10.6018/analesps.30.1.158201>
- Arce, R., Fariña, F., Seijo, D., & Novo, M. (2015). Assessing impression management with the MMPI-2 in child custody litigation. *Assessment*, 22(6), 769-777. <https://doi.org/10.1177/1073191114558111>
- Archer, R. P., Buffington-Vollum, J. K., Stredny, R. V., & Handel, R. W. (2006). A survey of psychological test use patterns among forensic psychologists. *Journal of Personality Assessment*, 87(1), 84-94. https://doi.org/10.1207/s15327752jpa8701_07
- Arias, E., Arce, R., Vázquez, M. J., & Marcos, V. (2020). Treatment efficacy on the cognitive competence of convicted intimate partner violence offenders. *Anales de Psicología/Annals of Psychology*, 36(3), 427-435. <https://doi.org/10.6018/analesps.428771>
- Baer, R. A., & Miller, J. (2002). Underreporting of psychopathology on the MMPI-2: A meta-analytic review. *Psychological Assessment*, 14(1), 16-26. <https://doi.org/10.1037/1040-3590.14.1.16>
- Bagby, R. M., & Marshall, M. B. (2004). Assessing underreporting response bias on the MMPI-2. *Assessment*, 11(2), 115-126. <https://doi.org/10.1177/1073191104265918>
- Basanta, J., Fariña, F., & Arce, R. (2018). Risk-need-responsivity model: Contrasting criminogenic and non-criminogenic needs in high and low risk juvenile offenders. *Children and Youth Services Review*, 85(1), 137-142. <https://doi.org/10.1016/j.childyouth.2017.12.024>
- Bonilla-Algovia, E., Rivas-Rivero, E. (2022). Papel del sexismo, los mitos románticos y los eventos vitales adversos en la violencia de género: un estudio con agresores de pareja [Influence of sexism, romantic myths and adverse life events on gender-based violence: A study with partner aggressors]. *Revista Iberoamericana de Psicología y Salud*, 13(2), 112-119. <https://doi.org/10.23923/j.rips.2022.02.057>
- Bonta, J., & Andrews, D. A. (2017). *The psychology of criminal conduct* (6th ed.). Routledge.
- Burchett, D. L., & Bagby, R. M. (2014). Multimethod assessment of response distortion: Integrating data from interviews, collateral records, and standardized assessment tools. In C. J. Hopwood & R. F. Bornstein (Eds.), *Multimethod clinical assessment* (pp. 345-378). Guilford Press.
- Butcher, J. N., Graham, J. R., Ben-Porath, Y. S., Tellegen, A., Dahlstrom, W. G., & Kaemmer, B. (2019). *MMPI-2: Inventario Multifásico de Personalidad de Minnesota-2* (4th ed.). TEA.
- Butcher, J. N., & Han, K. (1995). Development of the MMPI-2 scale to assess the presentation of self in a superlative manner: The S scale. In J. N. Butcher & C. D. Spielberger (Eds.), *Advance in personality assessment* (Vol. 10, pp. 25-50). LEA.
- Daffern, M., Papalia, N., Stevenson, E., Thomas, S. (2022). Common psychological treatments used to address criminal behavior. In C. Garofalo & J. J. Sijtsema (Eds.), *Clinical forensic psychology* (pp. 567-586). Palgrave Macmillan. https://doi.org/10.1007/978-3-030-80882-2_29
- Expósito-Álvarez, C., Lila, M., Gracia, E., & Martín-Fernández, M. (2021). Risk factors and treatment needs of batterer intervention program participants with substance abuse problems. *European Journal of Psychology Applied to Legal Context*, 13(2), 87-97. <https://doi.org/10.5093/ejpalc2021a9>
- Fandiño, R., Basanta, J., Sanmarco, J., Arce, R., & Fariña, F. (2021). Evaluation of the executive functioning and psychological adjustment of child to parent offenders: Epidemiology and quantification of harm. *Frontiers in Psychology*, 12, Article 616855. <https://doi.org/10.3389/fpsyg.2021.616855>
- Fariña, F., Arce, R., Vilarinho, M., & Novo, M. (2014). Assessment of the standard forensic procedure for the evaluation of psychological injury in intimate-partner violence. *Spanish Journal of Psychology*, 17, Article e32. <https://doi.org/10.1017/sjp.2014.30>
- Fariña, F., Redondo, L., Seijo, D., Novo, M., & Arce, R. (2017). A meta-analytic review of the MMPI validity scales and indexes to detect defensiveness in custody evaluations. *International Journal of Clinical and Health Psychology*, 17(2), 128-138. <https://doi.org/10.1016/j.ijchp.2017.02.002>
- Gancedo, Y., Fariña, F., Seijo, D., Vilarinho, M., & Arce, R. (2021). Reality monitoring: A meta-analytical review for forensic practice. *European Journal of Psychology Applied to Legal Context*, 13(2), 99-110. <https://doi.org/10.5093/ejpalc2021a10>
- Gancedo, Y., Sanmarco, J., Marcos, V., & Seijo, D. (2021). Estimating the efficacy of FPTSD Scale to report malingering of PTSD: A meta-analytic review. *Revista Iberoamericana de Psicología y Salud*, 12(1), 44-57. <https://doi.org/10.23923/j.rips.2021.01.044>
- Gancedo, Y., Sanmarco, J., Seijo, D., & Fariña, F. (2020). Malingering evaluation: A contrastive meta-analytic review of F and F-r scales. In M. Novo & D. Seijo (Eds.), *Psychology and Law: Research for practice* (pp. pp. 99-123). De Gruyter. <https://doi.org/10.2478/9788395669682-009>
- Gómez-Leal, R., Megías-Robles, A., Sánchez-López, M. T., & Fernández-Berrocal, P. (2021). Psychopathic traits and ability emotional intelligence in incarcerated males. *European Journal of Psychology Applied to Legal Context*, 13(2), 79-86. <https://doi.org/10.5093/ejpalc2021a8>
- Graham, J. R. (2011). *MMPI-2: Assessing personality and psychopathology* (5th ed.). Oxford University Press.
- Greene, R. L. (2011). *The MMPI-2/MMPI-2-RF: An interpretive manual* (3rd ed.). Allyn & Bacon.
- Jugl, I., Lösel, F., Bender, D., & King, S. (2021). Psychosocial prevention programs against radicalization and extremism: A meta-analysis of outcome evaluations. *European Journal of Psychology Applied to Legal Context*, 13(1), 37-46. <https://doi.org/10.5093/ejpalc2021a6>
- Ley Orgánica 1/1979, de 26 de septiembre, General Penitenciaria. (1979). *Boletín Oficial del Estado*, 239, 23180-23186. <https://www.boe.es/buscar/pdf/1979/BOE-A-1979-23708-consolidado.pdf>
- Ley Orgánica 1/2004, de 28 de diciembre, de Medidas de Protección Integral contra la Violencia de Género. *Boletín Oficial del Estado*, 313, 42166-42197. <https://www.boe.es/boe/dias/2004/12/29/pdfs/A42166-42197.pdf>
- Ley Orgánica 1/2015 de 30 de marzo, por la que se modifica la Ley Orgánica 10/1995, de 23 de noviembre, del Código Penal. *Boletín Oficial del*

- Estado, 77, 27061-27176. <https://www.boe.es/boe/dias/2015/03/31/pdfs/BOE-A-2015-3439.pdf>
- Ley Orgánica 3/2018, de 5 de diciembre, de Protección de Datos Personales y Garantía de los Derechos Digitales (2018). *Boletín Oficial del Estado*, 294, 119788-119857. <https://www.boe.es/boe/dias/2018/12/06/pdfs/BOE-A-2018-16673.pdf>
- Mayorga, E. G., Novo, M., Fariña, F., & Arce, R. (2020). Destrezas cognitivas en menores infractores, de protección y normalizados: un estudio de contraste [Cognitive skills in juvenile offenders, protection and normalized youngsters: A contrastive study]. *Revista Latinoamericana de Psicología*, 52, 160-168. <https://doi.org/10.14349/rlp.2020.v52.16>
- Molero, M. M., Martos, Á., Barragán, A. B., Pérez-Fuentes, M. C., & Gázquez, J. J. (2022). Anxiety and depression from cybervictimization in adolescents: A Metaanalysis and meta-regression study. *European Journal of Psychology Applied to Legal Context*, 14(1), 42 - 50. <https://doi.org/10.5093/ejpalc2022a5>
- Moral-Jiménez, M. V., & González-Sáez, M. E. (2020). Distorsiones cognitivas y estrategias de afrontamiento en jóvenes con dependencia emocional [Cognitive distortions and coping strategies in young people with emotional dependence]. *Revista Iberoamericana de Psicología y Salud*, 11(1), 15-30. <https://doi.org/10.23923/j.rips.2020.01.032>
- Nasaescu, E., Zych, I., Ortega-Ruiz, R., Farrington, D. P., & Llorent, V. J. (2020). Patrones longitudinales de las conductas antisociales en la adolescencia temprana: un análisis de clases y de transiciones latentes. *European Journal of Psychology Applied to Legal Context*, 12(2), 85-92. <https://doi.org/10.5093/ejpalc2020a10>
- Olson, C. L. (1976). On choosing a test statistic in multivariate analysis of variance. *Psychological Bulletin*, 83(4), 579-586. <https://psycnet.apa.org/doi/10.1037/0033-2909.83.4.579>
- Osuna, E., López-Martínez, M., Arce, R., & Vázquez, M. J. (2015). Analysis of response patterns on the MMPI-2 in psychiatric prison inmates. *International Journal of Clinical and Health Psychology*, 15(1), 29-36. <https://doi.org/10.1016/j.ijchp.2014.09.002>
- Paulhus, D. L. (1984). Two-component models of socially desirable responding. *Journal of Personality and Social Psychology*, 46(3), 598-609. <https://doi.org/10.1037/0022-3514.46.3.598>
- Podsakoff, P. M., MacKenzie, S. B., Lee, J.-Y., & Podsakoff, N. P. (2003). Common method bias in behavioral research: A critical review of the literature and recommended remedies. *Journal of Applied Psychology*, 88(5), 879-903. <https://doi.org/10.1037/0021-9010.88.5.879>
- Redondo, L., Fariña, F., Seijo, D., Novo, M., & Arce, R. (2019). A meta-analytical review of the responses in the MMPI-2/MMPI-2-RF clinical and restructured scales of parents in child custody dispute. *Anales de Psicología/Annals of Psychology*, 35(1) 156-165. <https://doi.org/10.6018/analesps.35.1.338381>
- Redondo, S., Padrón-Goya, F., & Martín, A. M. (2022). Offenders' narratives on criminal desistance while serving a prison sentence. *Victim and Offenders*, 17(3), 439-467. <https://doi.org/10.1080/15564886.2021.1910601>
- Rogers, R. (2018). Detection strategies for malingering and defensiveness. In R. Rogers & S. D. Bender (Eds.), *Clinical assessment of malingering and deception* (4th ed., pp. 18-41). Guilford Press.
- Rogers, R., Bagby, R. M., & Dickens, S. E. (1992). *The SIRS test manual*. Psychological Assessment Resources.
- Rogers, R., Sewell, K. W., Martin, M. A., & Vitacco, M. J. (2003). Detection of feigned mental disorders: A meta-analysis of the MMPI-2 and malingering. *Assessment*, 10(2), 160-177. <https://doi.org/10.1177/1073191103010002007>
- Rosenthal, R., & Rubin, D. B. (1982). A simple, general purpose display of magnitude of experimental effect. *Journal of Educational Psychology*, 74(2), 166-169. <https://doi.org/10.1037/0022-0663.74.2.166>
- Santirso, F. A., Lila, M., & Gracia, E. (2020). Motivational strategies, working alliance, and protherapeutic behaviors in batterer intervention programs: A randomized controlled trial. *European Journal of Psychology Applied to Legal Context*, 12(2), 77-84. <https://doi.org/10.5093/ejpalc2020a7>
- Sharf, A. J., Rogers, R., Williams, M. M., & Henry, S. A. (2017). The effectiveness of the MMPI-2-RF in detecting feigned mental disorders and cognitive deficits: A meta-analysis. *Journal of Psychopathology and Behavioral Assessment*, 39(3), 441-455. <https://doi.org/10.1007/s10862-017-9590-1>
- Stevens, J. (1986). *Applied multivariate statistics for social sciences*. LEA.
- Strong, D. R., Greene, R. L., Hope, C., Johnston, T., & Olesen, N. (1999). Taxometric analysis of impression management and self-deception on the MMPI-2 in child custody litigants. *Journal of Personality Assessment*, 73(1), 1-18. <https://doi.org/10.1207/S15327752JPA730101>
- Vilariño, M., Arce, R., & Fariña, F. (2013). Forensic-clinical interview: Reliability and validity for the evaluation of psychological injury. *European Journal of Psychology Applied to Legal Context*, 5(1), 1-21. <https://journals.copmadrid.org/ejpalc/archivos/articulo20180219092235.pdf>