

Psychoeducational Interventions in Children and Adolescents with Type-1 Diabetes: A Systematic Review

Bárbara Luque^{a,b}, Joaquín Villaécija^{a,b}, Rosario Castillo-Mayén^{a,b}, Esther Cuadrado^{a,b}, Sebastián Rubio^{a,b}, and Carmen Tabernero^{b,c,d}

^aUniversity of Córdoba, Spain; ^bMaimonides Biomedical Research Institute of Córdoba (IMIBIC), Spain; ^cSocial Psychology Department, University of Salamanca, Spain; ^dInstitute of Neurosciences of Castilla y León (INCYL), University of Salamanca, Spain

ARTICLE INFO

Article history:

Received 8 June 2021
Accepted 23 November 2021

Keywords:

Diabetes mellitus type 1
Education
Psychoeducation
Interventions
Children
Adolescents
Patient education
Systematic review

Palabras clave:

Diabetes tipo 1
Educación
Psicoeducación
Intervenciones
Niños
Adolescentes
Educación del paciente
Revisión sistemática

A B S T R A C T

The effectiveness of psychoeducational interventions in children and adolescents with type 1 diabetes is unclear. A systematic review was developed in accordance with PRISMA. Relevant databases (Pubmed, Cochrane, PsycINFO, and PsyARTICLES) were analyzed. Articles of the last decade with type 1 diabetes population between 6 and 18 years participating in psychoeducational interventions were the inclusion criteria. Twenty studies were reviewed, and improvements were found in glycosylated hemoglobin, diabetes knowledge, and psychosocial variables. The results support the positive effect of these interventions. The characteristics that seem to be behind the success of these interventions are the design appropriate to the characteristics of the population, the participation of psychologist and educators, the continuity of the program over time, and the use of digital tools and interaction strategies. Further studies need to be carried out and replicated in different groups of children and adolescents.

Las intervenciones psicoeducativas en los menores y adolescentes con diabetes tipo 1: una revisión sistemática

R E S U M E N

Hay dudas acerca de la efectividad de las intervenciones psicoeducativas en menores y adolescentes con diabetes tipo 1, motivo por el cual se realizó una revisión sistemática de acuerdo con el protocolo PRISMA. Se analizaron distintas bases de datos (Pubmed, Cochrane, PsycINFO y PsyARTICLES) con los siguientes criterios de inclusión: artículos de los últimos diez años, con población con diabetes tipo 1 de edades comprendidas entre los 6 y 18 años que hubieran participado en cualquier intervención psicoeducativa. Se revisaron 20 estudios y los resultados mostraron una mejora en la hemoglobina glicosilada, en el conocimiento de la enfermedad y en algunas variables psicosociales tras estas intervenciones. Las características que parecen estar detrás del éxito de estas intervenciones psicoeducativas son el diseño adecuado a las características de la población, la participación de profesionales de la psicología y de la educación, la continuidad del programa en el tiempo y el uso de herramientas digitales y otras estrategias de interacción. Se destaca la necesidad de realizar más estudios y que sean replicados en diferentes grupos de menores y adolescentes.

Chronic diseases are considered as noncommunicable diseases (NCDs) and cause many deaths each year. Specifically, 41 million people die each year from these diseases (71% of the deaths that occur worldwide in a year). Diabetes is included in this group of NCDs, along with cardiovascular diseases, cancer, and chronic respiratory diseases (World Health Organization [WHO, 2020]).

The main diseases are type 1 diabetes (T1D), type 2 diabetes (T2D), and gestational diabetes. T2D is the predominant type and occurs

mostly in adulthood. Although the body can produce insulin, it does not manage it correctly, and its origin is related to a deficit of physical activity and overweight. Gestational diabetes is a temporary condition during pregnancy that could complicate it, produced by an increase in blood glucose levels during this period (WHO, 2021). Finally, T1D, the focus of this study, normally appears at an early age, where the beta cells of the pancreas are attacked, losing its ability to produce insulin, the regulatory hormone of blood glucose levels (Spanish

Cite this article as: Luque, B., Villaécija, J., Castillo-Mayén, R., Cuadrado, E., Rubio, S., & Tabernero, C. (2022). Psychoeducational interventions in children and adolescents with type-1 diabetes: A systematic review. *Clínica y Salud*, 33(1), 35-43. <https://doi.org/10.5093/clysa2022a4>

Funding: This study was supported by the Spanish Ministry of Science, Innovation and University via a doctoral FPU grant to the second author (FPU18/04504), and was developed while a project was financially supported by the same public organization (PID2019-107304RB-I00), in which the main researchers are co-authors of the paper.

Correspondence: bluque@uco.es (B. Luque); carmen.tabernero@usal.es (C. Tabernero).

ISSN: 1130-5274/© 2022 Colegio Oficial de la Psicología de Madrid. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Diabetes Federation [FED, 2020]). Comorbidities of T1D include skin complications, neuropathy, foot problems, eye complications, DKA (ketoacidosis) and ketones, kidney disease, high blood pressure, and stroke (American Diabetes Association [ADA, 2020]).

It is estimated that currently 1.1 million children under 19 years of age have T1D worldwide, with 128,900 new cases diagnosed in children each year (International Diabetes Federation [IDF, 2019]). The prevalence differs between countries, although its frequency is increasing, especially in children under 5 years of age. Spain is the country with the highest incidence in southern Europe, with between 1,200 and 1,500 new cases diagnosed each year (Spanish Society of Pediatric Endocrinology [SSPE, 2019]).

The worrying data of this disease forces us to explore the scope of action that we have from education and psychology. The WHO has developed the Global Action Plan for the Prevention and Control of NCDs 2013-2020, where the education and empowerment of these patients has a fundamental role (WHO, 2013). In this sense, an educational intervention on diabetes, or diabetes self-management education (DSME), is a teaching-learning process about knowledge, tools, and practices for diabetes self-care that address the needs of the patient, to promote better health (Beck et al., 2017). These conventional educational programs are sometimes supplemented by psychosocial elements such as problem-solving, motivation, coping skills, stress management, counselling, communication skills, and behavioral therapy (Charalampopoulos et al., 2017; Murphy et al., 2006). These are known as psychoeducational interventions. In practice, both interventions are usually combined (Murphy et al., 2006).

However, the studies published on the effectiveness of educational or psychoeducational interventions in T1D show contradictory results. On the one hand, there is not enough evidence to justify that these programs are effective by themselves in children and young people with T1D, although there is evidence of their effectiveness when accompanied by other programs (Charalampopoulos et al., 2017; Murphy et al., 2006). On the other hand, positive results such as psychological, and educational benefits are found (Armour et al., 2005; Winkley et al., 2006), achieving better control of the disease and, consequently, a better quality of life for these patients. In addition, we cannot forget signs such as the benefit of the participation of psychologists, the rapid implementation in newly diagnosed children and the innovative strategies aimed at promoting patient participation (Charalampopoulos et al., 2017). Also, education appears to be most effective when integrated into routine care, when it encourages parental involvement and when adolescent self-efficacy (understood as the beliefs in one's own capabilities to achieve a goal in each situation) is promoted (Murphy et al., 2006; Wood & Bandura, 1989). Early interdisciplinary healthcare enhances the efficiency of disease management, and therefore, the quality of life of these patients (Urzealã et al., 2020). In adult population, which could serve as a background for another population group, one of the highlights is the use of mobile device App that could strengthen the perception of self-care by contributing to an increase in the information available about health education in diabetes, helping patients to control their glycated hemoglobin (HbA1c) (Bonoto et al., 2017).

The poor adaptation to the disease in the pediatric population (Bilbao-Cercós et al., 2014), understood as the degree of psychosocial adequacy of the subject's behavior, emotional state, and appraisal in relation to the disease (Portilla del Cañal & Jo, 1995), together with

the scarce scientific literature on the characteristics of educational and psychoeducational interventions in this population and the lack of an effective consolidated model, make it necessary to undertake this review to optimize the use of these interventions in children and adolescents with T1D in order to begin adaptation to the disease as soon as possible, since 90% of new diagnoses occur at this age (SSPE, 2019).

With this approach, the main aim of this study was to carry out an exploration on the approach of educational and psychoeducational interventions in children and adolescents (age range 6-18 years) with T1D, with two specific objectives: 1) to identify scientific evidence on the effectiveness of these interventions in this population and 2) to consider the methodological and educational strategies used in previous research to extract successful guidelines for designing future interventions.

Method

Study Design

Published scientific articles have been used to prepare the systematic review; therefore, ethical committee approval has not been necessary.

The study design was prepared in accordance with preferred reporting items for systematic reviews and meta-analyses (PRISMA) guidelines (Moher et al., 2009), as well as the instructions suggested by Cajal et al. (2020). To check the effectiveness of the interventions, we collected the psychosocial variables studied and, the HbA1c level as a biomedical indicator.

Inclusion Criteria

Scientific articles published in English or Spanish were searched, given that English is the language of scientific communication and Spanish is researchers' native language. To focus on current science and the latest educational trends, such as the use of technology and the rise of mHealth tools for self-care, only papers published during the last ten years (2010-2019) were used. Articles that did not allow access to the full text were eliminated. The country where the intervention was carried out was not considered as a reason for exclusion.

We defined the technical inclusion criteria by answering the PICO (acronym for patient, intervention, comparison and outcome) question, as shown in Table 1. Studies that did not differentiate between type 1 and type 2 diabetes mellitus populations have been excluded.

Search Strategy and Eligibility Criteria

The initial search was performed between October 2019 and February 2020. Four databases were used: Medline (PubMed), Cochrane Register of Controlled Trials (CENTRAL), PsycINFO, and PsycARTICLES. A search was also carried out in May 2020, before the final writing of the paper, to incorporate new studies published in this period.

The study population ranged between 6 and 18 years old. Studies were included if the mean age did not exceed 18 years old.

Table 1. PICO Strategy

Population	Children and adolescents (age range 6 to 18 years) with type 1 diabetes
Intervention	Educational or psychoeducational intervention
Control or comparator	Comparator that allows us to distinguish the effects (traditional control group, a pre-post design or an alternative intervention)
Outcome	The main outcome considered is the HbA1c level. Secondary outcomes: psychosocial variables (quality of life, diabetes knowledge, self-efficacy, and self-management).

Table 2. Search Strategies used in Databases

Medline (Pubmed)	((("type 1 diabetes"[Title/Abstract] OR t1d[Title/Abstract] OR "juvenile diabetes"[Title/Abstract] OR "insulin-dependent diabetes"[Title/Abstract] OR "type 1 diabetics"[Title/Abstract])) AND (education*[Title/Abstract] OR psychoeducation*[Title/Abstract] OR psycho-education*[Title/Abstract])) AND (children[Title/Abstract] OR adolescent*[Title/Abstract] OR youth[Title/Abstract] OR child[Title/Abstract] OR teenager*[Title/Abstract] OR "young people"[Title/Abstract])
Cochrane	"type 1 diabetes" OR t1d OR "juvenile diabetes" OR "insulin-dependent diabetes" OR "type 1 diabetics" in Title Abstract Keyword AND education OR psychoeducation OR psycho-education OR educational OR psychoeducational OR psycho-educational in Title Abstract Keyword AND children OR adolescent OR youth OR child OR teenager OR "young people" OR adolescents OR teenagers in Title Abstract Keyword
PsycINFO PsyARTICLES	ab("type 1 diabetes" OR t1d OR "juvenile diabetes" OR "insulin-dependent diabetes" OR "type 1 diabetics") AND ab(education* OR psychoeducation* OR psycho-education*) AND ab(children OR adolescent* OR youth OR child OR teenager* OR "young people")

Based on our PICO question, a combination of MESH terms (medical subject headings) was used: 1) "type 1 diabetes mellitus", "juvenile diabetes", "insulin-dependent diabetes", "type 1 diabetics"; 2) "education*", "psychoeducation*", "psycho-education*"; and 3) "children", "adolescent*", "youth", "child", "teenager*", "young people". These terms and their respective variants are used to search in title, abstract or keywords with the Boolean operators OR and AND, as shown in the detailed search in Table 2.

Strategies for the Selection of Studies and Analysis of the Results

The selection of articles was made independently by two researchers, and a third resolved any disagreements. Firstly, all publications found within the search criteria were transferred to the free version of the EndNote Clarivate Analytics platform and all the repeat publications were removed. A manual review was then necessary because some references did not match. The selection was first made by reading the title, then by reading the abstracts, and finally by selecting the studies to read in full.

title and 51 after the reading of abstract due to not meeting the inclusion criteria. In this way, 62 studies were selected to be read in full, of which only 20 were analyzed. The rest were discarded for the following justified reasons: other outcomes or type of study ($n = 12$), only the protocol was published ($n = 11$), pilot study ($n = 8$), invalid for inclusion criteria ($n = 5$), studies duplicated or intermediate balance ($n = 2$), no full text found ($n = 2$), or abstract of Congress type ($n = 2$). All the selected studies met the quality criteria evaluated using the Effective Public Health Practice Project (EPHPP) Quality Assessment Tool (Thomas et al., 2004). Figure 1 shows the flow diagram designed by PRISMA to analyze the different stages of study selection.

Characteristics of Selected Studies

Papers are summarized in Table 3. All studies included were published over the last decade (2010-2019), although some of them were developed in the years prior to the date of their publication; 75% of the studies were clinical trials. There were also cohort studies (10%), quasi-experimental studies (10%), and other not clearly specified (5%).

Studies from many locations resulted: Europe ($n = 9$), America ($n = 7$), Asia ($n = 2$), Africa ($n = 1$), and Oceania ($n = 1$). All studies selected involved a children or adolescents population with T1D. The sum of all studies results in 3,743 participants (53.52% females), of which 2,115 belonged to experimental groups with some type of educational or psychoeducational intervention. Study participants range from 24 to 675, and the mean of participants 187.2. The mean age groups ranged from 12.1 ± 1.1 years to 17.4 ± 2.4 years. Thirteen trials recruited adolescents (only over 11 years), while seven studies combined the child-youth population. HbA1c levels at baseline ranged from $7.8 \pm 1.1\%$ to $10.9 \pm 0.4\%$. For both age and HbA1c levels, we always prioritized having the mean at baseline, differentiating the group that performed the intervention and the control group, when this data is available. Other data are specified in the general characteristics Table 3.

Outcomes Found to Check the Effectiveness of the Intervention

The HbA1c level is the most used outcome to evaluate the impact of the educational or psychoeducational intervention; overall, 18 of the 20 selected studies used this variable, which was measured at baseline and at different times during and/or after the intervention. Among the psychosocial variables, the most studied was quality of life or health-related quality of life, which appeared in 12 studies, followed by self-management ($n = 5$), knowledge of diabetes ($n = 4$), and self-efficacy ($n = 4$).

Effectiveness of Interventions and Instruments Used

Of the 18 studies that measured HbA1c levels in their research, 11 provided significant improvements in this glycemic control variable. This improvement did not occur with the same analyses




Figure 1. Flow Diagram of Study Selection.

Results

Study Selection

A total of 1,040 potential scientific publications resulted from the designed search strategy: Medline ($n = 598$), Cochrane ($n = 329$), PsycINFO ($n = 107$), and PsyARTICLES ($n = 6$), although there were 869 final documents when duplicates were removed. Of these publications, 756 were excluded after the reading of

Table 3. Characteristics of Included Trials

First author (year) Country	Inclusion criteria	Intervention	Participants	Mean (SD) age (years)	Mean (SD) % HbA1c at baseline	Control group	Follow-up	Outcomes
Abolfotouh et al. (2011) Egypt	Age 12-20 years with T1D	Educational program. Four 120-minutes sessions, 1 per month in groups of 15 adolescents	IG: 121 CG: 122	14.6 ¹ (2.2)	10.5 ¹ (1.9)	Usual care	At baseline and 6 months	QoL; HbA1c
Altundang et al. (2016) Turkey	Age 12-14 years with T1D for at least 1 year	Group interaction (4-5 participants) in four 35-45-minutes sessions. Education Guide, and a consultation stage with family and peer interactions stages later	IG: 18 CG: 20	-	10.2 (2.4)	Usual care and Education Guide	At baseline and 3, 6 and 9 months	Self-Esteem; Social support; Diabetes knowledge
Brorsson et al. (2018) Sweden	Age 12-18 years with T1D > 12 months, HbA1c > 7.9%	GSD-Y, an empowerment-based, and person-centered reflection (2-hours each in 7 group sessions for 5 months)	IG: 37 CG: 32	14.8	8.4	Standard care	At baseline and 6 and 12 months	HbA1c; Self-perceived health; HRQoL; Family conflicts; Self-efficacy
Christie et al. (2016) UK	Age 8-16 years with T1D and a mean 12-month HbA1c of 8.5%	CASCADE. 4 module structured education which uses solution focused, and motivational approaches. Groups of 3-4 families. 1 session/month for 4 months	IG: 135/182 CG: 149/183	13.1 (2.1)	9.9 (1.5)	Standard care	At baseline and 12 and 24 months	HbA1c; QoL (Self-management; Intervention compliance; Emotional and behavioral adjustment)
Emiliana et al. (2019) Indonesia	Age 6-18 years with T1D	PRISMA. Educational tool with animated videos	IG: 31	-	-	Pre and Post type	At baseline and the end of intervention	Self-management; Level of compliance
García-Pérez et al. (2010) Spain	Age 11-18 years with T1D	Diabetes SME. Educational activities (interactive seminars and games to promote a healthier attitude) followed the recommendations of the ADA. In groups and individually. 8 days in a summer camp	IG: 34 CG: 23	13.8 (2.2)	8.6 (1.7)	Usual care	At baseline and at least 3 months after the camp	HbA1c; Diabetes knowledge; Anxiety; Psychological adaptation
Grey et al. (2013) USA	Age 11-14 years with T1D at least 6 months	TeenCope vs Managing Diabetes. Two Internet-based psycho-educational programs. One 30 minutes session/week for 5 weeks. TeenCope, a program based on social cognitive theory, used a graphic novel video format to model problematic social situations, and different coping skills to solve problems. Managing Diabetes was the CG	IG: 167 CG: 153	12.3 ¹ (1.1)	8.5 ¹ (1.4)	TeenCope vs. Managing Diabetes	At baseline and 3, 6, 12 and 18 months	Self-management; Level of compliance
Hawkes et al. (2019) USA	To be less than 18 years with T1D and for at least one diabetes autoantibody (we analyzed only 5-12 years and 12-18 years age groups)	T1Y1 program, a structured diabetes education. 10 hours throughout the year and families met with a nutritionist at 3 and 9 months after diagnosis	IG: 391 CG: 284	-	11.3	Standard Diabetes Care	At baseline and 6, 12, 18 and 24 months	HbA1c; QoL; and secondary (Coping, Self-efficacy; Social competence; Self-management; Family conflict)
Hood et al. (2018) USA	Age 14-18 years with T1D for at least 1 year and daily insulin dosing of at least 0.5 units/kg/day	STePS study. Distress and depression prevention program vs diabetes education program. Nine biweekly sessions lasting 90-120 minutes for 4.5 months	IG: 131 CG: 133	15.7 ¹ (1.1)	9.1 (2.0)	Resilience vs. education	At baseline and 4.5 (the end of intervention), 8, 12 and 16 months	DD; Diabetes self-management; and HbA1c
Iafusco et al. (2011) Italy	Age 10-18 years with T1D	A chat line intervention. Once a week (90-minutes) for at least 2 years consecutively. The topic of each session was vote by all participants at the beginning and concern about diabetes management, anxiety about future and social relationships	IG: 193 CG: 203	13.6 (2.7)	7.8 (1.1)	Non-chat group	At baseline and 1 and 2 years	QoL; HbA1c
Jaser et al. (2019) USA	Age 13-17 years with T1D for at least 6 months and HbA1c levels 8-12%	EDU (educational materials in mail every 2 weeks for 8 weeks with three-page packets and information about T1D) vs PA intervention. PA received the intervention reminders and the same materials that educational group, as well as components intended to induce PA	EDU: 60 PA: 60	14.8 ¹ (1.4)	9.2 ¹ (0.9)	EDU vs. PA	At baseline and 3 and 6 months	Adherence; HbA1c; QoL
Katz et al. (2014) USA	Age 8-16 years with T1D for at least 6 months	SC, CA+ (monthly outreach by the Care Ambassador via phone or email) and CA+Ultra (psychoeducational intervention that provided realistic expectations and problem-solving strategies).	SC: 51 CA+: 52 CA+Ultra: 50	12.9 ¹	8.4 ¹	SC, CA+ and CA+Ultra	At baseline and 1 and 2 years	HbA1c; QoL

Table 3. Characteristics of Included Trials (continued)

First author (year) Country	Inclusion criteria	Intervention	Participants	Mean (SD) age (years)	Mean (SD) % HbA1c at baseline	Control group	Follow-up	Outcomes
Mauri et al. (2017) Italy	Age 6-16 years with T1D	PED. 12-month structured project (educational workshops every three months based on active participation through play, and direct experiences) followed by an educational summer camp. Interdisciplinary team participation	IG: 24	12.1 (1.5)	8.8 (1.0)	Pre and Post type	At baseline (three-month before the project) and 1 year	HbA1c; Diabetes knowledge; Self-management; Wellbeing
Murphy et al. (2012) Australia	Adolescents with T1D for at least 1 year	FACTS. Six sessions (90 minutes monthly) incorporating skills training and family teamwork in 4-6 families/group	IG: 158 CG: 147	13.1* (1.9)	9.3* (1.9)	Usual care	At baseline and 6 and 18 months. HbA1c every 3 months	HbA1c; QoL; Well-being; Family responsibility
Petrovski et al. (2017) Macedonia	Age 14-23 years with T1D treated with an insulin pump and sensor for at least 6 months	Internet intervention for 3 years using Facebook and CareLink software. Intervention is the same as traditional but written reports and chats in Facebook. All patients received a standardized protocol of education about correct diabetes control	IG: 33 CG: 34	17.4 (2.4)	7.8 (1.8)	Standard care	At baseline and every 3 months during the study for a 3-year-period	HbA1c
Price et al. (2015) UK	Age 11-16 years with T1D for at least one year	KICK-OFF. An intensive education program. 5-day group education with interactive and practical activities	IG: 199 CG: 197	13.7 (1.4)	9.3 (1.7)	Usual care	At baseline and 6, 12 and 24 months	HbA1c; QoL; Diabetes Self-efficacy
Santiprabhob et al. (2012) Thailand	Patients older than 12 years with T1D	DSME and psychosocial support group (6-8 patients) sessions with problem-solving scenarios. 5-day camp	IG: 27	15.6 (2.1)	8.3 (1.8)	Pre and Post type	At baseline and 3, 6, 9 and 12 months	HbA1c; Diabetes knowledge; QoL; Self-care behavior
Verbeek et al. (2011) Netherlands	Age 11-17 years with T1D and level HbA1c > 9.0%	Psychoeducational program. 3 sessions (1.5 hours each) for patients and 1 for parents during a 3-month period	IG: 25	14.3 (1.7)	10.0 (0.7)	Pre and Post type	At baseline and 3 and 9 months	HbA1c
Wang et al. (2010) USA	Age 12-18 years with T1D for > 1 year and HbA1c ≥ than 9% on two consecutive visits	Motivational interviewing-based education (MI) vs structured diabetes education (SDE). Educators of MI were trained at a 2-day workshop	MI: 21 SDE: 23	MI: 15.3 (1.4) SDE: 15.6 (1.7)	MI: 10.9 (0.4) SDE: 11.1 (0.3)	MI vs SDE	At baseline and 3, 6 and 9 months	HbA1c; Depression; QoL; Self-care
Whittemore et al. (2016) USA	Age 11-14 years with T1D for at least 6 months	Internet Psychoeducational Program (Teens.Connect). Two components, TEENCOPE (5 interactive sessions) and Managing Diabetes (5 internet-based problem-solving lessons)	IG: 64 CG: 60	12.1 ¹ (1.1)	8.2 ¹ (1.4)	Planet D (Open access)	At baseline and 3 and 6 months	HbA1c; QoL; Self-care; Self-efficacy; Perceived stress; Depressive symptoms

Note. ADA = American Diabetes Association; CA = Care Ambassador; CA+ = Care Ambassador Plus; CA+Ultra = Care Ambassador Ultra; CASCADE = Child and Adolescent Structured Competencies Approach to Diabetes Education; CG = control group; DD = diabetes distress; DSME = Diabetes Self-Management Education; EDU = Education; FACTS = Families and Adolescents Communication and Teamwork Study diabetes education program; GSD-Y = Guided Self-Determination-Young; HbA1c = glycosylated hemoglobin; HRQoL = health-related quality of life; IG = intervention group; KICK-OFF = Kids in Control of Food; MI = Motivational Interviewing-based education; PA = positive affect; QoL = quality of life; PED = Pediatric Education for Diabetes; SC = Standard Care; SDE = Structured Diabetes Education; SE = Structured Education; SME = Self-Management Education; StePS = Supporting Teens Problem Solving; T1D = type 1 diabetes; T1Y1 = Type 1 Year 1 program.

¹Undifferentiated data between groups.

but can be classified into three groups: produced after comparison with the levels of the experimental group itself prior to the study ($n = 3$), due to a comparison between the experimental group and the control group ($n = 5$), or some other improvement due to the interaction of another biomedical or psychosocial variable ($n = 3$). Regarding psychosocial variables, positive effects were also found: the quality of life measure showed improvements in 5 studies, diabetes knowledge showed improvement in all trials in which it is used ($n = 4$), self-efficacy in 2 studies, and self-management in 1. In addition, other less-used variables also showed improvements, such as self-esteem, social support, distress diseases, or perceived stress. For the measurement of psychosocial variables included in our study, where improvements were found, the instruments used are described. For the quality of life, the Diabetes Quality of Life for Youth Inventory (DQOLY) (Iafusco et al., 2011), with a Cronbach alpha for its three factors of .85 in life satisfaction, .83 in disease impact, and .82 in disease-related worries in the original questionnaire (Ingersoll & Marrero, 1991). In the Arabic version of DQOLY (Abolfotouh et al., 2011), Cronbach alpha was .83; in the Pediatric Quality of Life Inventory (PedsQL) (Grey et al., 2013), Cronbach alpha was .87 in the studied sample; in the Pediatric Quality of Life Inventory Type 1 Diabetes Module (PedsQL-D) (Jaser et al., 2019; Price et al., 2016), Cronbach alpha was .71 in the original scale (Varni

et al., 2003); and in the Generic Quality of Life (PedsQL-G) (Price et al., 2016), Cronbach alpha was .88 in the original (Varni et al., 2001). For diabetes knowledge, a test was created by the researchers in one study (Altundag & Bayat, 2016); a questionnaire based on a Spanish adult population was adapted for children and adolescents (García-Pérez et al., 2010), with a Cronbach alpha of .63, and .87 in the original (DISK) (Bueno et al., 1993); some questions were used from the Diabetes Knowledge Questionnaire [Questionario sulla conoscenza del diabete] from the Italian Diabetes Education Study Group (GISED), with a Cronbach alpha of .60; and another non-specific instrument (Santiprabhob et al., 2012). For self-efficacy, a diabetes-specific subscale of self-efficacy for the Diabetes Scale was used, and Cronbach alpha was .88 in the studied sample (Grey et al., 2013). A questionnaire was designed (Abolfotouh et al., 2011), similar to the design by McCaul et al. (1987). For self-management, a self-management questionnaire was used with a reliability test of .91 (Emiliana et al., 2019).

General Features of the Intervention

All the studies selected according to the inclusion criteria had a methodological design that allowed to measure the effects of the educational or psychoeducational intervention. Three methods were

found: intervention group versus control group ($n = 13$), pre- and post-type ($n = 4$), and comparison between different interventions ($n = 3$); 85% of the studies included psychosocial variables to control the effectiveness of the interventions, while the rest (15%) took the biomedical variable HbA1c as the only outcome.

The educational perspective shows a heterogeneous implementation of multiple strategies and resources for the development of these interventions. Programs range from a more traditional model of diabetes education to more innovative models that work on psychosocial variables. The involvement of information and communication technologies should be noted: chat line (Iafusco et al., 2011), Facebook (Petrovski & Zivkovic, 2017), animated videos (Emiliana et al., 2019), or other forms of Internet intervention (Grey et al., 2013; Whittemore et al., 2016). Even so, most interventions (75%)

did not make use of digital resources: 4 of the 5 studies that rely on digital resources to carry out their interventions showed significant improvements in HbA1c levels. Various psychoeducational strategies were explicitly found: empowerment-based (Borsson et al., 2019), motivational approaches (Christie et al., 2016; Wang et al., 2010), distress program (Hood et al., 2018), positive affect (PA) (Jaser et al., 2019), role-playing (Grey et al., 2013; Katz et al., 2014; Mauri et al., 2017; Price et al., 2016; Santiprabhob et al., 2012; Whittemore et al., 2016). Six studies were found in which psychologists participated in different ways: in the intervention (García-Pérez et al., 2010; Iafusco et al., 2011; Mauri et al., 2017; Santiprabhob et al., 2012), in a pilot study before intervention (Christie et al., 2016), and training diabetes educator (Wang et al., 2010).

All the details of the methods are described in the Intervention-

Table 4. Main Contributions of the Studies Analyzed: Conclusions and Outcomes

Study	Main contribution
Abolfotouh et al. (2011)	C: Educational intervention may protect QoL and glycemic control from worsening over time. O: Deterioration QoL in IG was significantly less severe than in the CG; knowledge, adherence, self-efficacy and HbA1c improved ($p < .001$).
Altundang et al. (2016)	C: Training and peer interaction were found to be effective in adapting to the disease. O: There was a decrease in HbA1c levels and an increase in self-esteem ($p < .001$) and social support ($p < .05$), and significant increase in knowledge levels ($p < .001$) in IG. No changes in CG.
Borsson et al. (2018)	C: An intervention with GSD-Y may have an effect on glycemic control. The content may serve as a model for person-centered care. O: HbA1c with an analysis between boys and girls separately and adjusted for family conflicts, boys after 12 months improved ($p = .019$).
Christie et al. (2016)	C: Significant challenges in the delivery of a SE intervention using psychological techniques to enhance engagement and behavior change. O: HbA1c levels did not improve in children and adolescents with poor control.
Emiliana et al. (2019)	C: The use of animated videos in SME could improve self-management and children's compliance in the management of diet, physical act, treatment, stress management and blood glucose control. O: PRISMA education had significant effects on self-management and level of compliance ($p < .05$).
García-Pérez et al. (2010)	C; O: No relevant changes in diabetes knowledge, anxiety or psychological adaptation were found after the psychoeducational intervention in the summer camp. The adaptation to the school environment was the only significantly improvement.
Grey et al. (2013)	C: Internet interventions improved results. Combining both diabetes management education and behavioral interventions is more effective than only one. O: After 18 months who completed both interventions had lower HbA1c ($p = .04$) and higher QoL ($p = .02$) and self-efficacy ($p = .03$), among others, compared with who did only one.
Hawkes et al. (2019)	C: SE and support in the first year after diagnosis can improve short-term outcomes, although the effect did not persist after training. O: HbA1c was significantly lower in the T1Y1 group at 6 ($p < .001$), 12 ($p < .001$) and 18 ($p < .01$) months. No effect at 24 months.
Hood et al. (2018)	C: Intervention before your psychological symptoms begin can prevent DD. STePS represents a promising prevention program. O: Intervention was associated with reduction in DD between groups ($p < .05$) and stable glycemic control, resilience characteristics and depressive symptoms in the 1 year post-treatment.
Iafusco et al. (2011)	C: A chat line is an effective tool to the diabetes team that could help to improve diabetes compliance. O: QoL improved in the intervention group ($p = .001$). Decreased HbA1c levels in the intervention group ($p < .001$) although no difference was observed between groups ($p = .056$).
Jaser et al. (2019)	O: A positive psychology intervention had initial significant positive effects on coping and quality of life. A more intensive or longer-lasting intervention may be needed to sustain these effects and to improve adherence and glycemic control. O: No significant effects were found for glycemic control. PA intervention group improved quality of life ($p = .022$) and disengagement coping at 3 months ($p = .018$), but not at 6 months.
Katz et al. (2014)	C: The psychoeducational intervention was effective in maintaining or improving HbA1c. O: There were no differences in HbA1c across treatment groups although more youth in the psychoeducation group with suboptimal baseline HbA1c $\geq 8\%$ maintained or improved their HbA1c.
Mauri et al. (2017)	C: A model where a pediatric diabetologist cooperates with an adult diabetologist seems be a solution to the transitional gap. O: HbA1c improved significantly ($p < .05$), knowledge of self-monitoring of blood glucose and the diabetes adapted nutrition ($p < .001$) and the presence of adequate behaviors ($p < .001$).
Murphy et al. (2012)	C: Attendance at group education sessions in clinics was very poor. More personalized educational approaches may be required. O: No significant difference in HbA1c at 18 months.
Petrovski et al. (2017)	C: Social networks like Facebook can help improve glycemic control using insulin pump therapy. O: Both groups improved HbA1c levels. Significant improvement ($p < .05$) was found in favor of the Internet group.
Price et al. (2015)	C; O: HbA1c levels showed no significant improvement. Improvements in QoL score levels at 6 months.
Santiprabhob et al. (2012)	C: The effect of diabetes camp on glycemic control is controversial. O: HbA1c levels and the QoL did not improve statistically significant post-camp. Knowledge 12 months post-camp improved ($p < .001$).
Verbeek et al. (2011)	C: A psychoeducational program can be beneficial to improve HbA1c levels. O: HbA1c levels decreased after 9 months ($p = .08$). In a subgroup of 15 patients showed a significant reduction at 9 months follow-up.
Wang et al. (2010)	C: SDE is found to be effective in improving metabolic control. O: Over the follow-up, the SDE group had a lower adjusted mean HbA1c than MI group ($p = .03$). There were not differences in psychosocial measures.
Whittemore et al. (2016)	C: Teens need frequent reminders to increase their participation in psychoeducational programs. O: After 6 months there were no significant differences in HbA1c, QoL or other outcomes between groups. Teen. Connect group had lower perceived stress over time ($p < .01$).

Note. C = conclusions; CG = control group; DD = diabetes distress; GSD-Y = Guided Self-Determination-Young; HbA1c = glycosylated hemoglobin; IG = intervention group; MI = Motivational Interviewing-based education; O = outcomes; PA = positive affect; QoL = quality of life; SDE = Structured Diabetes Education; SE = Structured Education; SME = Self-Management Education; STePS = Supporting Teens Problem Solving; T1Y1 = Type 1 Year program.

column of [Table 3](#), where specific characteristics of the programs are outlined. Regarding the main contributions made by each study and their most outstanding results, they are presented in [Table 4](#).

Discussion

The aim of this systematic review was to carry out an exploration to find scientific evidence of the positive impact of educational or psychoeducational interventions, and the strategies used for its development to improve disease control in children and adolescents with T1D. HbA1c levels were the most common outcome measure. Although the improvement of HbA1c level is important, from an education and psychology viewpoint other motivational and psychosocial variables that allow us to achieve greater control of the disease are also relevant. Perhaps, the most important is self-management, the goal of any educational intervention on diabetes.

Contrasting Positions on the Effectiveness of Educational or Psychoeducational Interventions

There is insufficient evidence to recommend the use of specific psychoeducational interventions for children and adolescents with T1D, based on a review of UK trials ([Charalampopoulos et al., 2017](#)). Even so, as this same study confesses, meta-analyses in the USA found that psychoeducational interventions can improve HbA1c levels by up to half a percentage point, in addition to other psychological and educational benefits ([Armour et al., 2005](#); [Winkley et al., 2006](#)). Another study argues that although no evidence of significant improvements in HbA1c levels was found in their meta-analysis for the adolescent population after structured education, there was evidence for the adult population ([Liu et al., 2020](#)), though a reduction in vascular complications was found in the adult population with T1D ([Menezes et al., 2016](#)).

The need to continue studying theoretical approaches and methods that can be applied, including the strategies of these programs, remains latent, since education can lead to better control of diabetes ([Jenhani et al., 2005](#); [Pals et al., 2020](#)). If one method is effective for one population, could it be effective for another one if it is adapted appropriately? The heterogeneity of the interventions makes it impossible for us to adopt a single stance because of diverse implementations. Strategies, methods, and tools have worked to provide indications that could be addressed and combined in future interventions.

Featured Strategies, Methods, and Tools

Firstly, the use of digital tools seems to favor the positive effect. Of the five analyzed studies that are supported by digital resources and tools, all five present a significant improvement in some of their analyzed variables: HbA1c levels ([Grey et al., 2013](#); [lafusco et al., 2011](#); [Petrovski & Zivkovic, 2017](#)), self-management ([Emiliana et al., 2019](#)), self-efficacy ([Grey et al., 2013](#)), and quality of life ([Grey et al., 2013](#); [lafusco et al., 2011](#)). Specifically, animated videos could improve self-management of the disease ([Emiliana et al., 2019](#)); internet interventions combining diabetes management education and behavioral interventions ([Grey et al., 2013](#)); a chat line ([lafusco et al., 2011](#)); and the use of social networks as a platform to deliver these interventions to improve glycemic control using insulin pump therapy ([Petrovski & Zivkovic, 2017](#)). The use of information and communication technology (ICT) creates expectations, yet to be discovered, with great potential for the treatment of chronic diseases such as diabetes ([Rhee et al., 2020](#)).

This strengthens the idea that using applications could help improve HbA1c control and strengthen the perception of self-care and safety in diabetic patients ([Bonoto et al., 2017](#)). Although the

use of video games is also recommended as a potential tool for educational interventions, it needs to be specifically designed for that age group and framed within the theoretical foundations of health psychology ([DeShazo et al., 2010](#)). In this sense, the use of these tools could facilitate an intervention based on behavioral models to promote the development of self-efficacy judgements in this population, according to the social cognitive theory formulated by [Bandura \(1997\)](#).

On the other hand, peer interaction is a strategy that can benefit the achievement of objectives in an educational setting. Specifically, training and peer interaction could be effective in adapting to the disease ([Altundag & Bayat, 2016](#)). Peer-based interventions show some promise, although there are not many studies ([Kazemi et al., 2016](#)). For the transition stage, the cooperation of an adolescent and an adult, both with diabetes disease, can be an effective option for a progressive transition of care ([Mauri et al., 2017](#)).

The continuity over time of the interventions also seems to be a key factor in their success. Positive effects have been found with a positive psychology intervention ([Jaser et al., 2019](#)), or with a structured education program ([Hawkes et al., 2019](#)), but the effects did not continue after training. A more intensive or longer-lasting intervention may be needed to sustain these effects ([Jaser et al., 2019](#)). Families who re-visited the web portal after one year obtained better glycemic control ([Hanberger et al., 2013](#)). It is probable that young people need frequent reminders to increase their participation in psychoeducational interventions ([Whittemore et al., 2016](#)).

It is also important to highlight that different methodologies and various approaches achieved positive results with their interventions, e.g., self-management courses ([Johnson et al., 2019](#)), structured diabetes education ([Wang et al., 2010](#)), psychoeducational programs ([Katz et al., 2014](#); [Verbeek et al., 2011](#)), and positive psychology interventions ([Jaser et al., 2019](#)), among others. The heterogeneity of the methods used, and their effectiveness, leads us to believe that adaptation to the specific context in which it will be applied is what is truly important ([Charalampopoulos et al., 2017](#)), rather than a particular educational program ([Murphy et al., 2006](#)).

Finally, the implementation of these intervention programs is mostly carried out by medical specialists and nurses in nutrition and diabetes care. Even so, the participation of physicians or multidisciplinary staff has shown effect ([Menezes et al., 2016](#)), and the involvement of psychologists was one of the differences with the successful USA programs ([Charalampopoulos et al., 2017](#)). We believe that the participation of psychologists is fundamental for successful psychosocial and motivational variables, and the participation of experts in education is fundamental for the pedagogical and methodological approach of interventions where the teaching-learning process has a primary role.

Limitations and Future Approaches

In accordance with the main aim initially stated, we consider that the development of the research has been correct and has allowed an exploration of the educational and psychoeducational interventions developed in children and adolescents with T1D. In relation to the first specific objective, some of the limitations encountered do not allow us to affirm the effectiveness of these interventions with absolute certainty, although there are indications of their usefulness. In relation to the second aim, techniques and methods have been found that appear to be effective in the development of these interventions.

It is possible that some very specific studies have been left out of the selection due to not being included under these criteria. In this way, this study could be extended to other databases, languages, and other criteria to enrich its results. The wide age range chosen could also be a limitation, although due to the limited scientific literature on this topic, we have decided not to adjust it further. In

this sense, as research in this field increases, future studies could consider establishing greater differentiation in the stage of the life cycle at which the interventions are targeted taking into account the classification in the stages of human development (e.g., [Papalia & Martorell, 2017](#)).

Regarding the content, the heterogeneity of the interventions analyzed led us to the decision not to carry out a meta-analysis since there are many variables that could modify an objective result. The main opportunity that it offers us is the possibility of taking the conclusions of the analyzed interventions as a reference, and the successful guidelines discussed, for the design of future interventions. Regarding patients' age, one of the limitations of the study lies in the fact that the articles reviewed are heterogeneous in terms of the age at which they direct their psychoeducational interventions, therefore the conclusions should be taken with caution.

Conclusions

The results obtained and subsequent discussion lead us to believe that educational and psychoeducational interventions have the potential to improve management of the disease and other psychosocial variables in children and adolescents with T1D. We cannot affirm that these interventions are always effective by themselves. It is necessary for more studies to be carried out with a larger population, and that these studies be replicated with the same design in different groups of children and adolescents, considering the characteristics of this population and their interests, to determine their efficacy.

In summary, as a basic premise, we believe that an effective intervention must be designed in accordance with the setting and the population in which it is going to be implemented. Even though there is no evidence of a successful valid model, we have found potential indicators that could serve as guidelines for future interventions and further research on them. Psychologists and educators should be involved in the design and supervise these interventions. Professionals trained in teaching-learning processes, together with specialist medical staff and diabetes educator (generally nurses), create a balance of knowledge and that would allow the continuing study of the effectiveness of these programs. In addition, the correct choice of resources and educational methods depends on the participation of these educators and psychologists. In this regard, there are two issues to consider continuing in the line of research: a resource, digital tools in their different aspects (animated videos, portal web, social networks, Apps, etc.) and a strategy, interaction (peers' group, people's experiences, role-play, solving problems, etc.). Both aspects seem that they could facilitate a positive impact on diabetes education and therefore, the control of the disease. Finally, another important factor to consider is the continuity of the programs over time to extend their effectiveness.

References

References marked with an asterisk indicate studies included in the meta-analysis.

- *Abolfotouh, M. A., Kamal, M. M., El-Bourgy, M. D., & Mohamed, S. G. (2011). Quality of life and glycemic control in adolescents with type 1 diabetes and the impact of an education intervention. *International Journal of General Medicine*, 4, 141-152. <https://doi.org/10.2147/IJGM.S16951>
- *Altundag, S., & Bayat, M. (2016). Peer interaction and group education for adaptation to disease in adolescents with type 1 diabetes mellitus. *Pakistan Journal of Medical Sciences*, 32(4), 1010-1014. <https://doi.org/10.12669/pjms.324.9809>
- American Diabetes Association (ADA, 2020). *Diabetes type 1 complications*. <https://www.diabetes.org/diabetes/complications>
- Armour, T. A., Norris, S. L., Jack, L., Zhang, X., & Fisher L. (2005). The effectiveness of family interventions in people with diabetes mellitus: A systematic review. *Diabetic Medicine*, 22(10), 1295-1305. <https://doi.org/10.1111/j.1464-5491.2005.01618.x>

- Bandura, A. (1997). *Self-efficacy: The exercise of control*. W. H. Freeman.
- Beck, J., Greenwood, D. A., Blanton, L., Bollinger, S. T., Butcher, M. K., Condon, J. E., Cypress, M., Faulkner, P., Fischl, A. H., Francis, T., Kolb, L. E., Lavin-Tompkins, J. M., McLeod, J., Maryniuk, M., Mensing, C., Orzeck, E. A., Pope, D. D., Pulizzi, J. L., Reed, A. A., ... Wang, J. (2017). 2017 National standards for diabetes self-management education and support. *Diabetes Care*, 40(10), 1409-1419. <https://doi.org/10.2337/dci17-0025>
- Bilbao-Cercós, A., Beniel-Navarro, D., Pérez-Marín, M., Montoya-Castilla, I., Alcón-Sáez, J. J., & Prado-Gascó, V. J. (2014). El autoconcepto y la adaptación a la enfermedad en pacientes diabéticos pediátricos. *Clinica y Salud*, 25(1), 57-65. <https://doi.org/10.5093/cl2014a5>
- Bonoto, B. C., de Araújo, V. E., Godói, I. P., de Lemos, L. L., Godman, B., Bennie, M., Diniz, L. M., & Junior, A. A. (2017). Efficacy of mobile Apps to support the care of patients with diabetes mellitus: A systematic review and meta-analysis of randomized controlled trials. *JMIR mHealth and eHealth*, 5(3), e4. <https://doi.org/10.2196/mhealth.6309>
- *Brorsson, A. L., Leksell, J., Andersson-Franko, M., & Lindholm-Olinder, A. (2019). A person-centered education for adolescents with type 1 diabetes-A randomized controlled trial. *Pediatric Diabetes*, 20(7), 986-996. <https://doi.org/10.1111/pedi.12888>
- Bueno, J. M., Marco, M. D., Leal, A., Orozco, D., & Mira, J. J. (1993). An evaluation study of a scale of diabetological education in primary care. *Atención Primaria*, 11(7), 344-348.
- Cajal, B., Jiménez, R., Gervilla, E., & Montañó, J. J. (2020). Doing a systematic review in health sciences. *Clinica y Salud*, 31(2), 77-83. <https://doi.org/10.5093/cl2020a15>
- Charalampopoulos, D., Hesketh, K. R., Amin, R., Paes, V. M., Viner, R. M., & Stephenson, T. (2017). Psycho-educational interventions for children and young people with type 1 diabetes in the UK: How effective are they? A systematic review and meta-analysis. *PLoS One*, 12(6), e0179685. <https://doi.org/10.1371/journal.pone.0179685>
- *Christie, D., Thompson, R., Sawtell, M., Allen, E., Cairns, J., Smith, F., Jamieson, E., Hargreaves, K., Ingold, A., Brooks, L., Wiggins, M., Oliver, S., Jones, R., Elbourne, D., Santos, A., Wong, I. C. K., O'Neil, S., Strange, V., Hindmarsh, P., ... Viner, R. M. (2016). Effectiveness of a structured educational intervention using psychological delivery methods in children and adolescents with poorly controlled type 1 diabetes: A cluster-randomized controlled trial of the CASCADE intervention. *BMJ Open Diabetes Research & Care*, 4(1), e000165. <https://doi.org/10.1136/bmjdr-2015-000165>
- DeShazo, J., Harris, L., & Pratt, W. (2010). Effective intervention or child's play? A review of video games for diabetes education. *Diabetes Technology & Therapeutics*, 12(10), 815-822. <https://doi.org/10.1089/dia.2010.0030>
- *Emiliana, P., Agustini, N., Allenidekania, & Rustina, Y. (2019). A preliminary study on "PRISMA" education in improving self-management and level of compliance in children with type-1 diabetes mellitus. *Comprehensive Child and Adolescent Nursing*, 42(sup1), 115-121. <https://doi.org/10.1080/24694193.2019.1578432>
- *García-Pérez, L., Perestelo-Pérez, L., Serrano-Aguilar, P., & Del Mar Trujillo-Martín, M. (2010). Effectiveness of a psychoeducative intervention in a summer camp for children with type 1 diabetes mellitus. *The Diabetes Educator*, 36(2), 310-317. <https://doi.org/10.1177/0145721710361784>
- *Grey, M., Whittemore, R., Jeon, S., Murphy, K., Faulkner, M. S., Delamater, A., & TeenCope Study Group (2013). Internet psycho-education programs improve outcomes in youth with type 1 diabetes. *Diabetes Care*, 36(9), 2475-2482. <https://doi.org/10.2337/dci12-2199>
- Hanberger, L., Ludvigsson, J., & Nordfeldt, S. (2013). Use of a web 2.0 portal to improve education and communication in young patients with families: randomized controlled trial. *Journal of Medical Internet Research*, 15(8), e175. <https://doi.org/10.2196/jmir.2425>
- *Hawkes, C. P., Willi, S. M., & Murphy, K. M. (2019). A structured 1-year education program for children with newly diagnosed type 1 diabetes improves early glycemic control. *Pediatric Diabetes*, 20(4), 460-467. <https://doi.org/10.1111/pedi.12849>
- *Hood, K. K., Iturralde, E., Rausch, J., & Weissberg-Benchell, J. (2018). Preventing diabetes distress in adolescents with type 1 diabetes: Results 1 year after participation in the STePS Program. *Diabetes Care*, 41(8), 1623-1630. <https://doi.org/10.2337/dci17-2556>
- *Iafusco, D., Galderisi, A., Nocerino, I., Cocca, A., Zuccotti, G., Prisco, F., & Scaramuzza, A. (2011). Chat line for adolescents with type 1 diabetes: A useful tool to improve coping with diabetes: a 2-year follow-up study. *Diabetes Technology & Therapeutics*, 13(5), 551-555. <https://doi.org/10.1089/dia.2010.0188>
- Ingersoll, G. M., & Marrero, D. G. (1991). A modified quality-of-life measure for youths: Psychometric properties. *The Diabetes Educator*, 17(2), 114-118. <https://doi.org/10.1177/014572179101700219>
- International Diabetes Federation (IDF, 2019). *IDF diabetes atlas* (9th ed.). <https://www.diabetesatlas.org>
- *Jaser, S. S., Whittemore, R., Choi, L., Nwosu, S., & Russell, W. E. (2019). Randomized trial of a positive psychology intervention for adolescents with type 1 diabetes. *Journal of Pediatric Psychology*, 44(5), 620-629. <https://doi.org/10.1093/jpepsy/jsz006>
- Jenhani, M., Gaha, K., Nabouli, R., Ghedira, A., & Abdelaziz, A. (2005). Effectiveness of patient education on glycemic control in insulin treated patients in general practice. *Diabetes & Metabolism*, 31(4), 376-381. [https://doi.org/10.1016/S1262-3636\(07\)70207-0](https://doi.org/10.1016/S1262-3636(07)70207-0)

- Johnson, B., Norman, P., Sanders, T., Elliott, J., Whitehead, V., Campbell, F., Hammond, P., Ajjan, R., & Heller, S. (2019). Working with insulin, carbohydrates, ketones and exercise to manage diabetes (WICKED): Evaluation of a self-management course for young people with type 1 diabetes. *Diabetic Medicine*, 36(11), 1460-1467. <https://doi.org/10.1111/dme.14077>
- *Katz, M. L., Volkening, L. K., Butler, D. A., Anderson, B. J., & Laffel, L. M. (2014). Family-based psychoeducation and Care Ambassador intervention to improve glycemic control in youth with type 1 diabetes: A randomized trial. *Pediatric Diabetes*, 15(2), 142-150. <https://doi.org/10.1111/pedi.12065>
- Kazemi, S., Parvizy, S., Atlasi, R., & Baradaran, H. R. (2016). Evaluating the effectiveness of peer-based intervention in managing type I diabetes mellitus among children and adolescents: A systematic review. *Medical Journal of the Islamic Republic of Iran*, 30, 442.
- Liu, F., Guan, Y., Li, X., Xie, Y., He, J., Zhou, Z. G., & Li, L. (2020). Different effects of structured education on glycemic control and psychological outcomes in adolescent and adult patients with type 1 diabetes: A systematic review and meta-analysis. *International Journal of Endocrinology*, 2020, e9796019. <https://doi.org/10.1155/2020/9796019>
- *Mauri, A., Schmidt, S., Soser, V., Sambataro, M., Nollino, L., Fabris, F., Corò, A., Scantamburlo, A., Marcon, M. L., Cazziola-Merlotto, M., Ciani, T., Tessarin, M., & Paccagnella, A. (2017). A structured therapeutic education program for children and adolescents with type 1 diabetes: An analysis of the efficacy of the "Pediatric Education for Diabetes" project. *Minerva Pediatrics*, 73(2), 159-166. <https://doi.org/10.23736/S0026-4946.17.04634-5>
- McCaul, K. D., Glasgow, R. E., & Schafer, L. C. (1987). Diabetes regimen behaviors. Predicting adherence. *Medical Care*, 25(9), 868-881. <https://doi.org/10.1097/00005650-198709000-00005>
- Menezes, M. M., Lopes, C. T., & Nogueira, L. (2016). Impact of educational interventions in reducing diabetic complications: A systematic review. *Revista Brasileira de Enfermagem*, 69(4), 773-784. <https://doi.org/10.1590/0034-7167.2016690422i>
- Moher, D., Liberati, A., Tetzlaff, J., Altman, D. G., & PRISMA Group (2009). Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Medicine*, 6(7), e1000097. <https://doi.org/10.1371/journal.pmed.1000097>
- Murphy, H. R., Rayman, G., & Skinner, T. C. (2006). Psycho-educational interventions for children and young people with type 1 diabetes. *Diabetic Medicine*, 23(9), 935-943. <https://doi.org/10.1111/j.1464-5491.2006.01816.x>
- *Murphy, H. R., Wadham, C., Hassler-Hurst, J., Rayman, G., Skinner, T. C., & Families and Adolescents Communication and Teamwork Study (FACTS) Group (2012). Randomized trial of a diabetes self-management education and family teamwork intervention in adolescents with type 1 diabetes. *Diabetic Medicine*, 29(8), e249-e254. <https://doi.org/10.1111/j.1464-5491.2012.03683.x>
- Pals, R. A. S., Skinner, T., Velasco, E. R., & Grabowski, D. (2020). The role of theories in interventions targeting preteens with type 1 diabetes: A critical literature review. *Child: Care, Health and Development*, 46(2), 155-174. <https://doi.org/10.1111/cch.12730>
- Papalia, D., & Martorell, G. (2017). *Desarrollo humano* [Human development]. McGraw Hill Education.
- *Petrovski, G., & Zivkovic, M. (2017). Impact of Facebook on glucose control in type 1 diabetes: A three-year cohort study. *JMIR diabetes*, 2(1), e9. <https://doi.org/10.2196/diabetes.7693>
- Portilla del Cañal, L., & Jo, A. S. (1995). Construcción y validación de un cuestionario para evaluar la respuesta adaptativa a la enfermedad en pacientes diabéticos. *Revista Cubana de Psicología*, 12(3), 193-201.
- *Price, K. J., Knowles, J. A., Fox, M., Wales, J. K., Heller, S., Eiser, C., Freeman, J. V., & KICK-OFF Study group (2016). Effectiveness of the kids in control of food (KICK-OFF) structured education course for 11-16 years old with type 1 diabetes. *Diabetic Medicine*, 33(2), 192-203. <https://doi.org/10.1111/dme.12881>
- Rhee, S. Y., Kim, C., Shin, D. W., & Steinhubl, S. R. (2020). Present and future of digital health in diabetes and metabolic disease. *Diabetes & Metabolism Journal*, 44(6), 819-827. <https://doi.org/10.4093/dmj.2020.0088>
- *Santiprabhob, J., Kiattisakthavee, P., Likitmaskul, S., Chaichanwattanakul, K., Wekawanich, J., Dumrongphol, H., Sriwijitkamol, A., Peerapatdit, T., & Nitiyanant, W. (2012). Glycemic control, quality of life and self-care behavior among adolescents with type 1 diabetes who attended a diabetes camp. *The Southeast Asian Journal of Tropical Medicine and Public Health*, 43(1), 172-184.
- Spanish Diabetes Federation (SDF, 2020). *Types of diabetes*. https://fedesp.es/diabetes/tipos/#Diabetes_tipo_1
- Spanish Society of Pediatric Endocrinology (SSPE, 2019). *What you should know about diabetes in the pediatric age*. https://www.mscbs.gob.es/organizacion/sns/planCalidadSNS/pdf/excelencia/cuidadospalativos-diabetes/DIABETES/Lo_que_debes_saber_sobre_la_diabetes_en_la_edad_pediatica.pdf
- Thomas, B. H., Ciliska, D., Dobbins, M., & Micucci, S. (2004). A process for systematically reviewing the literature: Providing the research evidence for public health nursing interventions. *Worldviews on Evidence-Based Nursing*, 1(3), 176-184. <https://doi.org/10.1111/j.1524-475X.2004.04006.x>
- Urzealã, C., Bota, A., Teodorescu, S., Vlăiculescu, M., Baker, J. S., & The Kidscreen Group Europe (2020). Quality of life in Romanian children with type 1 diabetes: A cross-sectional survey using an interdisciplinary healthcare intervention. *Healthcare*, 8(4), 382. <https://doi.org/10.3390/healthcare8040382>
- Varni, J. W., Burwinkle, T. M., Jacobs, J. R., Gottschalk, M., Kaufman, F., & Jones, K. L. (2003). The PedsQL in type 1 and type 2 diabetes: Reliability and validity of the Pediatric Quality of Life Inventory generic core scales and type 1 diabetes module. *Diabetes Care*, 26(3), 631-637. <https://doi.org/10.2337/diacare.26.3.631>
- Varni, J. W., Seid, M., & Kurtin, P. S. (2001). PedsQL 4.0: Reliability and validity of the Pediatric Quality of Life Inventory version 4.0 generic core scales in healthy and patient populations. *Medical Care*, 39(8), 800-812. <https://doi.org/10.1097/00005650-200108000-00006>
- *Verbeek, S., Vos, R. C., Mul, D., & Houdijk, M. E. (2011). The influence of an educational program on the HbA(1c)-level of adolescents with type 1 diabetes mellitus: A retrospective study. *Journal of Pediatric Endocrinology & Metabolism*, 24(1-2), 15-19. <https://doi.org/10.1515/jpem.2011.104>
- *Wang, Y. C., Stewart, S. M., Mackenzie, M., Nakonezny, P. A., Edwards, D., & White, P. C. (2010). A randomized controlled trial comparing motivational interviewing in education to structured diabetes education in teens with type 1 diabetes. *Diabetes Care*, 33(8), 1741-1743. <https://doi.org/10.2337/dc10-0019>
- *Whittemore, R., Liberti, L. S., Jeon, S., Chao, A., Minges, K. E., Murphy, K., & Grey, M. (2016). Efficacy and implementation of an Internet psychoeducational program for teens with type 1 diabetes. *Pediatric Diabetes*, 17(8), 567-575. <https://doi.org/10.1111/pedi.12338>
- Winkley, K., Ismail, K., Landau, S., & Eisler, I. (2006). Psychological interventions to improve glycaemic control in patients with type 1 diabetes: Systematic review and meta-analysis of randomised controlled trials. *BMJ (Clinical research ed.)*, 333(7558), 65. <https://doi.org/10.1136/bmj.38874.652569.55>
- Wood, R., & Bandura, A. (1989). Impact of conceptions of ability on self-regulatory mechanisms and complex decision making. *Journal of Personality and Social Psychology*, 56(3), 407-415. <https://doi.org/10.1037//0022-3514.56.3.407>
- World Health Organization (WHO, 2013). *Global action plan for the prevention and control of NCDs 2013-2020*. <https://www.who.int/publications/i/item/9789241506236>
- World Health Organization (WHO, 2020). *Noncommunicable diseases*. <https://www.who.int/news-room/fact-sheets/detail/noncommunicable-diseases>
- World Health Organization (WHO, 2021). *Diabetes*. <https://www.who.int/news-room/fact-sheets/detail/diabetes>

