



The European Journal of Psychology Applied to Legal Context

<https://journals.comadrid.org/ejpalc>



Motivational Strategies, Working Alliance, and Protherapeutic Behaviors in Batterer Intervention Programs: A Randomized Controlled Trial

Faraj A. Santirso, Marisol Lila, and Enrique Gracia

University of Valencia, Spain

ARTICLE INFO

Article history:

Received 12 April 2020
Accepted 11 May 2020

Keywords:

Batterer intervention program
Motivational strategies
Working alliance
Protherapeutic behaviors
Randomized controlled trial

ABSTRACT

Motivational strategies are among the most promising approaches to improve the effectiveness of batterer intervention programs (BIPs). An individualized motivational plan (IMP) is one of these motivational strategies. The present study aimed to explore whether adding an IMP to a standard BIP improved the participant-facilitator working alliance and participants' protherapeutic behaviors. To this end a randomized controlled trial was conducted. One hundred fifty-three men convicted of intimate partner violence were randomly assigned to either a standard BIP (control condition, $n = 79$) or a standard BIP plus IMP (experimental condition, $n = 74$). Working alliance (i.e., general working alliance, agreement, and bond) was assessed with the Working Alliance Inventory-Observer, short version. Protherapeutic behaviors (i.e., assumption of responsibility, participant role behavior, and group value) were assessed with the Observational Coding of Protherapeutic Group Behavior. Both working alliance and protherapeutic behaviors were assessed by an external observer early and late in intervention. Our results showed that both general working alliance and agreement and bond, were significantly higher in the standard BIP plus IMP intervention condition, both early and late in intervention. All protherapeutic behaviors were significantly higher in the standard BIP plus IMP early in intervention, and also late in intervention for assumption of responsibility and group value. Our findings have important practical implications as our results clearly showed that a motivational strategy tool such as the IMP improves key intervention processes (i.e., working alliance and protherapeutic behaviors) in BIPs, therefore increasing their effectiveness.

Las estrategias motivacionales, la alianza de trabajo y la conducta proterapéutica en los programas de intervención con maltratadores: un ensayo clínico aleatorizado

RESUMEN

Las estrategias motivacionales se encuentran entre los enfoques más prometedores para mejorar la eficacia de los programas de intervención con maltratadores. El plan motivacional individualizado (PMI) es una de estas estrategias motivacionales. El presente estudio tiene como objetivo explorar si añadir un plan motivacional individualizado a un programa estándar de intervención con maltratadores mejora la alianza de trabajo facilitador-participante y la conducta proterapéutica de los participantes. Para ello se realizó un ensayo clínico aleatorizado. Ciento cincuenta y tres hombres condenados por violencia de género fueron asignados aleatoriamente bien a un programa estándar de intervención con maltratadores (condición control, $n = 79$) o bien a un programa estándar de intervención con maltratadores más PMI (condición experimental, $n = 74$). La alianza de trabajo (i.e., alianza general, acuerdo y vínculo) se evaluó con la versión breve del *Working Alliance Inventory-Observer*. Las conductas proterapéuticas (i.e., asunción de responsabilidad, rol conductual del participante y valoración del grupo) fueron evaluadas con el *Observational Coding of Protherapeutic Group Behavior*. Tanto la alianza de trabajo como las conductas proterapéuticas fueron evaluadas por un observador externo al principio y al final de la intervención. Los resultados mostraron que tanto la alianza de trabajo general como el acuerdo y el vínculo fueron significativamente mayores en la condición experimental, tanto al principio como al final de la intervención. La expresión de todas las conductas proterapéuticas al inicio de la intervención fue significativamente mayor en la condición experimental, así como al final de la intervención para la asunción de responsabilidad y la valoración del grupo. Los resultados tienen importantes implicaciones prácticas, puesto que muestran con claridad que una estrategia motivacional como el PMI mejora procesos clave de la intervención con maltratadores (i.e., la alianza de trabajo y las conductas proterapéuticas), mejorando por lo tanto la efectividad de estos programas.

Palabras clave:

Programas de intervención con maltratadores
Estrategias motivacionales
Alianza de trabajo
Conductas proterapéuticas
Ensayo clínico aleatorizado

Cite this article as: Santirso, F. A., Lila, M., & Gracia, E. (2020). Motivational strategies, working alliance, and protherapeutic behaviors in batterer intervention programs: A randomized controlled trial. *The European Journal of Psychology Applied to Legal Context*, 12(2), 77-84. <https://doi.org/10.5093/ejpalc2020a7>

Funding: This research was supported by the Spanish Ministry of Health, Consumption, and Social Services, National Drug Plan (PND2018/021). Faraj A. Santirso was supported by the FPU Program of the Spanish Ministry of Education, Culture and Sport (grant number FPU15/00864).

Correspondence: Marisol.Lila@uv.es (M. Lila).

ISSN: 1889-1861/© 2020 Colegio Oficial de la Psicología de Madrid. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Several meta-analyses conducted to evaluate the effectiveness of batterer intervention programs (BIPs) have indicated that the effect size of these interventions on reducing recidivism tends to be small (Arce et al., 2020; Babcock et al., 2004; Cheng et al., 2019; Eckhardt et al., 2013; Feder & Wilson, 2005; Smedslund et al., 2011). This limited success of BIPs in reducing recidivism has been attributed to different factors such as low treatment compliance and high dropout rates (Bennett et al., 2007; Daly & Pelowski, 2000; Jewell & Wormith, 2010; Lila et al., 2019; Olver et al., 2011), lack of motivation for change (Carbajosa, Catalá-Miñana, Lila, & Gracia, 2017; Carbajosa, Catalá-Miñana, Lila, Gracia, et al., 2017; Crane et al., 2015; Eckhardt et al., 2008; Zalmanowitz et al., 2013), problems in building the working alliance (Cadsy et al., 1996; DiGiuseppe et al., 1994; Murphy & Baxter, 1997; Taft et al., 2004), and poor engagement in program activities (Musser et al., 2008; Taft et al., 2003).

Motivational strategies are one of the approaches to overcome these limitations and, consequently, to improve the effectiveness of BIPs (Babcock et al., 2016; Feder & Wilson, 2005; Santirso et al., 2020). Among the motivational strategies included in BIPs are, for example, interventions based on the 'stage of change' approach (Alexander et al., 2010), retention techniques (Mbilinyi et al., 2011; Musser et al., 2008; Taft et al., 2001), or motivational interviewing (Alexander et al., 2010; Crane & Eckhardt, 2013; Kistenmacher & Weiss, 2008; Lila et al., 2018; Mbilinyi et al., 2011; Murphy et al., 2017; Musser et al., 2008; Scott et al., 2011; Woodin & O'Leary, 2010). A growing body of research reports promise results regarding the effectiveness of BIPs incorporating motivational strategies, in terms of decreasing intimate partner violence recidivism and levels of dropout (Alexander et al., 2010; Crane & Eckhardt, 2013; Lila et al., 2018; Mbilinyi et al., 2011; Murphy et al., 2017; Scott et al., 2011; Taft et al., 2001; Woodin & O'Leary, 2013) and increasing intervention attendance, homework compliance, and stage of change level (Crane & Eckhardt, 2013; Kistenmacher & Weiss, 2008; Lila et al., 2018; Murphy et al., 2017; Musser et al., 2008; Taft et al., 2001).

Motivational strategies aim to build a strong facilitator-participant working alliance that helps to reduce participants' intervention resistance and increase motivation for change (Stuart et al., 2007). Also, these strategies are likely to promote protherapeutic behaviors among participants, thus leading to an improved group climate in BIPs (Brown & O'Leary, 2000; Musser et al., 2008; Rondeau et al., 2001; Semiatin et al., 2013; Taft et al., 2003; Taft et al., 2004).

The working alliance, according to Bordin (1979), consists of three components: facilitator-participant agreement on the intervention goals, participants' acceptance of and collaboration with the tasks proposed by facilitators to address participants' problems, and facilitator-participant emotional bond. The working alliance has been related to physical and psychological IPV reduction. For example, using observers' scores, Brown and O'Leary (2000) found that the working alliance in the first sessions of a voluntary couple intervention on IPV was associated with significant reductions of physical and psychological violence at the end of the intervention. In a court-mandated group intervention to prevent IPV, Taft et al. (2003) found that the therapeutic alliance reported by the therapist was also related with significant reductions of physical and psychological violence at six-month follow-up.

As for protherapeutic behaviors in the context of BIPs, Semiatin et al. (2013) defined them in terms of participants' verbalizations indicating: a) acknowledgment of personal responsibility and the need for personal change to avoid future violent behavior; b) participants' role behavior facilitating positive changes of other group members; and c) participants' verbalizations indicating a positive perception of both the group and the intervention program. These protherapeutic behaviors have been related to a significant reduction of physical and psychological IPV reported by victims at six-month follow-up (Semiatin et al., 2013).

However, despite the importance of these key intervention processes, little research has been conducted to assess the effectiveness of motivational strategies on increasing participant-facilitator working alliance and participants' protherapeutic behavior in BIPs.

The Present Study

The main aim of the present study was to examine whether adding motivational strategies to a standard BIP increases participant-facilitator working alliance and participants' protherapeutic behaviors. To this end, a randomized controlled trial was conducted, comparing a standard BIP (control group) with an experimental condition in which an individualized motivational plan was added to the standard BIP as a motivational strategy (IMP; Lila et al., 2018; Romero-Martínez et al., 2019). The IMP is based on several approaches aimed at increasing intervention compliance in BIPs that have proved their effectiveness in different settings: motivational interviewing (Miller & Rollnick, 2002), stages of change approach (Prochaska & DiClemente, 1982; Prochaska et al., 1992), solution-focused brief therapy (DeShazer & Berg, 1997), and the Good Lives model (Langlands et al., 2009; Ward, 2002). The main components of the IMP are: (1) five individual motivational interviews, of which three were conducted during the evaluation phase to reduce participants' resistance to the intervention and establish participants' personal goals, one was conducted in the middle of the program to monitor their progress, and the last one was conducted around the end of the program to assess their achievements; (2) three group sessions, where participants share their personal goals, explain their progress to the group, and receive feedback, support, and advice from facilitators and other group participants (these group sessions are conducted at the beginning, middle, and end of the program); (3) facilitators follow-up and reinforcement of participants' goals in every weekly group session throughout the intervention; and (4) retention techniques, such as phone calls when participants miss a group session. The IMP also implies that facilitators adopt an empathic and motivational attitude throughout the intervention, creating a climate of acceptance and using confrontation only when it becomes absolutely necessary.

As far as we know, the only previous experimental study analyzing whether motivational strategies increase working alliance and protherapeutic behaviors in BIP participants was conducted by Musser et al. (2008). In their study, groups receiving two motivational interviewing sessions as a pretreatment intervention were compared to control groups (i.e., intake as usual, without motivational interviewing). These researchers observed in the experimental group an increase in protherapeutic behaviors (i.e., verbalizations of responsibility for abusive behavior and group intervention usefulness) and higher working alliance late in intervention rated by therapists.

The present study pulls ahead previous research in two relevant issues, one related to the implementation of motivational strategies and the other related to the assessment of both protherapeutic behaviors and working alliance. First, in our study the motivational strategy was delivered throughout the BIP and not only as pretreatment intervention. We expected that the higher the exposure to motivational strategies the more positive the effect on the working alliance and protherapeutic behaviors. Second, we used a systematic observational methodology conducted by external observers to assess both protherapeutic behaviors and the working alliance. This methodological approach addressed impression management and social desirability issues when using offenders' self-reports, as well as potential biases in facilitators' self-reports or observations (Bennett, 2007; Gracia et al., 2015; Juarros-Basterrechea et al., 2018; Santirso et al., 2018; Weber et al., 2019).




Figure 1. Flow Diagram.

Method

Participants

The sample consisted of 153 males convicted of intimate partner violence against women and court-mandated to a community-based BIP. Offenders had been sentenced to less than a two-year term in prison, did not have previous criminal records, and their sentence was suspended on the condition that they attended the intervention. Eligibility criteria for this study were the following: (a) men over 18 years of age, (b) who had no severe psychological disorder, (c) had no severe substance abuse problems, and (d) had signed an informed consent form. Mean age was 40.73 years ($SD = 11.99$, range: 18-78). The sample consisted primarily of Spanish (71.7%, $n = 110$), Latin American ($n = 17$, 11.2%), European (other than Spanish, $n = 13$, 8.6%), African ($n = 11$, 7.2%), and Asian ($n = 2$, 1.3%) males. Regarding educational level, 6.4% had no education, 51% had primary education, 31.5% had secondary education, and 11.1% had university education. As for marital status, 33.3% were single ($n = 51$), 32.7% were divorced ($n = 50$), 24.2% were married or in a relationship ($n = 37$), and 9.8% were separated ($n = 15$). On average, annual family household income was between €6,000 and €12,000. About half of the participants were unemployed 45.1% ($n = 69$) at the time of the initial assessment.

Intervention Conditions

Standard batterer intervention program (SBIP). This condition consisted of 35 weekly group sessions of a standard cognitive-

behavioral intervention. It was divided into six modules with the following main aims: a) first module: to build a climate of trust and establish norms for the group to function; b) second module: to introduce the IPV basic concepts and address attribution of responsibility; c) third module: to train in cognitive emotion management techniques and cognitive restructuring; d) fourth module: to develop awareness of IPV consequences on victims, empathy, and positive communication skills in intimate relationships; e) fifth module: to discuss sexist attitudes, gender roles, and gender equality; and f) sixth module: to consolidate learning objectives and prevent relapse. Several techniques were applied during the SBIP (i.e., group dynamics, role-playing, monitored exercises, and training in cognitive restructuring or emotion management skills). Closed-ended groups in both conditions ranged from 10 to 12 men per group and were led by two facilitators.

Standard batterer intervention program plus individualized motivational plan (SBIP + IMP). The experimental condition consisted of the same standard cognitive-behavioral intervention with the addition of the IMP (see description above).

Facilitators training and intervention adherence. Facilitators were psychologists with at least two years' experience in BIPs. They received approximately 25 hours of training in their respective intervention condition. Facilitators were blind to the intervention condition. Each pair of facilitators intervened exclusively on one intervention condition. Facilitators for each condition were supervised independently once every two weeks. Supervision sessions focused on adherence to treatment protocol, group management, participants' progress, and preparation of future sessions. To ensure the content of

and adherence to the protocol, written intervention manuals for each condition were used (Lila et al., 2018).

Randomization. Participants assessed for eligibility came through the penitentiary system ($N = 181$). Twenty-eight men were excluded, mainly for not attending the first meeting (see Figure 1). A random number generator was used to allocate participants to the SBIP or the SBIP + IMP condition. Fourteen intervention groups were established, seven for the SBIP + IMP condition ($n = 74$) and seven for the SBIP condition ($n = 79$). Figure 1 provides the description of participant flow from recruitment to study completion.

Procedure

Participants were clearly told that refusing to participate in the study would not affect their legal situation and would have no legal consequences. Confidentiality was assured, with the sole exception of situations that could pose a risk to participants or other people. Participants who agreed to take part in this study completed a written consent form and were randomly assigned to the SBIP + IMP or SBIP condition. Four independent trained graduate research assistants, who were blind to the objectives and hypotheses of the study, coded

videotaped sessions. Working alliance and protherapeutic group behaviors were assessed twice: early in intervention (sessions 3-7) and late in intervention (sessions 24-28). Raters previously underwent training in which they assessed the same recorded session separately until they reached an acceptable level of agreement (i.e., not differing by more than one point on each assessed item). Two-hour recorded intervention sessions were divided into 24 five-minute intervals. Each participant received an average rating across session intervals on working alliance and protherapeutic group behaviors.

Measures

Working Alliance Inventory-Observer short version (WAI-O-S; Tichenor & Hill, 1989; Spanish version by Santirso et al., 2018). This observational scale assesses both general working alliance, and two components of working alliance (i.e., agreement and bond). The WAI-O-S contains 12 items (e.g., "There is a mutual liking between participant and facilitator", "There is agreement on what is important for the participant to work on"). Raters responded on a 7-point Likert-type scale ranging from 0 (*conclusive evidence against*) to 7 (*conclusive evidence in favor*). The scale had adequate internal

Table 1. Baseline Characteristics of Participants in Each Intervention Condition ($N = 153$)

Variables	SBIP + IMP ¹ ($n = 74$)		SBIP ² ($n = 79$)		t	χ^2
	$M(SD)$	%	$M(SD)$	%		
Age	39.39 (11.66)		41.99 (12.23)		1.34 (ns)	
Income ³	4.31 (2.53)		4.00 (2.12)		- 0.83 (ns)	
Origin						0.44 (ns)
Spain		69.9		73.4		
Latin American		12.3		10.1		
Europe (excluding Spain)		9.6		7.6		
Africa		6.8		7.6		
Asia		1.4		1.3		
Education						4.46 (ns)
No education		2.7		10.1		
Primary		50.0		51.9		
Secondary		36.5		26.6		
University		10.8		11.4		
Marital status						4.59 (ns)
Married or with partner		18.9		29.1		
Single		36.5		30.4		
Separated		6.8		12.7		
Divorced		37.8		27.8		
Unemployed		50.0		40.5		1.39 (ns)

Note. ¹SBIP + IMP = standardized batterer intervention program plus individualized motivational plan; ²SBIP = standardized batterer intervention program; ³annual income: 1 = < €1,800; 2 = €1,800-€3,600; 3 = €3,600-€6,000; 4 = €6,000-€12,000; 5 = €12,000-18,000; 6 = €18,000-€24,000; 7 = €24,000-€30,000; 8 = €30,000-€36,000; 9 = €36,000-€60,000; 10 = €60,000-€90,000; 11 = €90,000-€120,000; and 12 = > €120,000.

Table 2. Descriptive Statistics and 2 x 2 Repeated Measures ANOVAs for Completers Sample ($n = 104$)

	Early in Intervention		Late in Intervention		Time	Group	Time * Group			
	SBIP + IMP ¹ ($n = 54$)	SBIP ² ($n = 50$)	SBIP + IMP ¹ ($n = 54$)	SBIP ² ($n = 50$)						
	$M(SD)$	$M(SD)$	$M(SD)$	$M(SD)$						
General working alliance	4.07 (0.15)	3.92 (0.29)	4.13 (0.21)	4.04 (0.12)	15.30***	.13	12.50***	.11	1.51	.02
Agreement	4.09 (0.16)	3.93 (0.34)	4.15 (0.24)	4.06 (0.16)	15.65***	.13	9.77**	.09	2.11	.02
Bond	4.04 (0.15)	3.91 (0.23)	4.10 (0.19)	4.00 (0.08)	12.49***	.11	16.63***	.14	0.53	.01
Responsibility for abuse	3.01 (0.11)	2.87 (0.26)	3.03 (0.11)	2.98 (0.10)	12.81***	.11	13.91***	.12	5.23*	.05
Participant role behavior	3.01 (0.10)	2.89 (0.26)	3.01 (0.04)	2.99 (0.09)	7.00***	.06	10.63***	.09	7.31**	.07
Group value	3.00 (0.01)	2.94 (0.14)	3.06 (0.14)	2.99 (0.03)	13.20***	.12	20.85***	.17	0.04	.01

Note. SBIP + IMP¹ = standardized batterer intervention program plus individualized motivational plan; SBIP² = standardized batterer intervention program.

* $p \leq .05$, ** $p \leq .01$, *** $p \leq .001$.

consistency in this study with Cronbach's alpha equal to .97 and .92 for early and late in intervention measures, respectively. In a previous study with a sample of male IPV offenders, results showed an excellent level of inter-rater agreement and significant correlations with other indicators of intervention effectiveness (e.g., stage of change, motivation to change, and protherapeutic group behaviors; see Santirso et al., 2018).




Figure 2. Working Alliance Scores Early in Intervention and Late in intervention in SBIP+IMP group (represented by triangles) and SBIP group (represented by squares). a) General working alliance; b) Agreement subscale; c) Bond subscale. * Effect of group (SBIP + IMP or SBIP) ($p < 0.05$).

Observational coding of protherapeutic group behavior (Semiatin et al., 2013). This observational tool assesses protherapeutic behaviors of group participants through their verbalizations. It is a 3-item measure of the following protherapeutic behaviors: (a) responsibility for abuse—participants' verbalizations related to assuming vs. denying responsibility for their abusive actions, consequences of these actions, and the need for a personal change to avoid committing abusive acts in the future; (b) participant role behavior—interpersonal behaviors within the group that promote or

hinder change by other participants; the coding system addressed four types of participant role behavior along two axes, confirmation vs. confrontation and negative progress vs. positive progress; and (c) group value—participant verbalizations related to the perceived value of the group and the intervention in general. Raters assessed each of the protherapeutic behaviors on a 5-point Likert-type scale ranging from 1 (*conclusive evidence against*) to 5 (*conclusive evidence in favor*). The effect of different raters assessing the same participants was evaluated. Pooled reliability correlation (r) across the three coded variables for averaged session ratings was .53, $n = 30$, $p = .011$.

Statistical Analyses

To analyze whether participants who received SBIP + IMP and participants who received SBIP were equivalent at the time of allocation, chi-square and independent t -tests were conducted for categorical and continuous variables, respectively. To assess 'time' and 'group' differences in working alliance and protherapeutic behavior, we carried out repeated measures ANOVAs, with working alliance and protherapeutic group behaviors (early and late in intervention) as between-subject factors, and 'intervention group' (SBIP + IMP or SBIP) as the within-subject factor. When a factor was significant in previous ANOVAs, Bonferroni tests were performed. All statistical analyses were conducted using SPSS 26.0, and two-tailed tests with p set to .05 were considered as significant.

Results

Baseline Characteristics

Table 1 contains information on sociodemographic characteristics by treatment condition. The results showed that randomization was satisfactory. No pretreatment characteristics significantly distinguished SBIP + IMP and SBIP groups.

Observational Ratings of Working Alliance and Protherapeutic Group Behaviors

There was a significant effect of 'time' on general working alliance, and on the agreement and bond subscales, $F(1, 102) = 15.30$, $p = .0001$, $\eta^2 = .13$; $F(1, 102) = 15.65$, $p = .0001$, $\eta^2 = .13$; $F(1, 102) = 12.49$, $p = .001$, $\eta^2 = .11$, respectively, with higher scores later in intervention than early in intervention in the total sample (see Table 2). A significant effect of 'group' was also found on general working alliance, and on agreement and bond subscales, $F(1, 102) = 12.50$, $p = .001$, $\eta^2 = .11$; $F(1, 102) = 9.77$, $p = .002$, $\eta^2 = .09$; $F(1, 102) = 16.63$, $p = .0001$, $\eta^2 = .14$, respectively. Size effects were moderate to large (Cohen, 1988). Participants who received SBIP + IMP intervention showed higher general working alliance than those who received SBIP intervention, regardless of intervention moment (see Table 2 and Figure 2). No other significant effects were found on working alliance.

Additionally, a significant effect of 'time' was found on responsibility for abuse, participant role behavior, and group value scores, $F(1, 102) = 12.81$, $p = .001$, $\eta^2 = .11$; $F(1, 102) = 7.00$, $p = .009$, $\eta^2 = .06$; $F(1, 102) = 13.20$, $p = .0001$, $\eta^2 = .12$, respectively, with higher scores late in intervention compared to early in intervention in the total sample (see Table 2). Moreover, a significant 'group' effect was found on responsibility for abuse, participant role behavior and group value scores, $F(1, 102) = 13.92$, $p = .0001$, $\eta^2 = .12$; $F(1, 102) = 10.63$, $p = .002$, $\eta^2 = .09$, $F(1, 102) = 20.85$, $p = .0001$, $\eta^2 = .17$, respectively, with participants who received SBIP + IMP having higher scores (see Figure 3). Size effects were moderate to large (Cohen, 1988). Furthermore, a significant 'time * group' effect was

found on responsibility for abuse and participant role behavior scores, $F(1, 102) = 5.23, p = .02, \eta^2 = .05$; $F(1, 102) = 7.31, p = .008, \eta^2 = .07$, respectively (see Table 2 and Figure 3). Specifically, participants who received SBIP + IMP intervention showed higher responsibility for abuse early ($p = .001$) and late in intervention ($p = .013$) than those who received SBIP. Additionally, participants who received SBIP + IMP condition had significantly higher participant role behavior scores ($p = .002$) and tended to show higher participant role behaviors scores late in treatment than those who received SBIP only ($p = .067$). Finally, responsibility for abuse and participant role behaviors significantly increased throughout intervention in participants who received SBIP (for all, $p < .0001$), but not in those in SBIP + IMP condition (for all, $p > .35$).




Figure 3. Protherapeutic Group Behavior Early in Intervention and Late in Intervention in SBIP+IMP group (represented by triangles) and SBIP group (represented by squares). a) Responsibility for abuse; b) Participant role behavior; c) Group value.

* Effect of group (SBIP + IMP or SBIP) ($p < 0.05$). * Effect of 'time*group' ($p < 0.05$). (t) Effect of 'time*group' ($p = 0.067$).

Discussion

Working alliance and protherapeutic behaviors are key intervention processes to improve BIP effectiveness. Motivational strategies can contribute to build a strong working alliance and promote protherapeutic behaviors. However, little rigorous research has been conducted to assess the effectiveness of motivational strategies to increase working alliance and participants' protherapeutic behavior in BIPs. The randomized controlled trial conducted in this study showed that adding motivational strategies to a standard BIP increases participant-facilitator working alliance and participants' protherapeutic behaviors.

Our results showed that both general working alliance and scores on the agreement and bond subscales increased significantly over the course of the intervention in both intervention conditions, and was significantly higher in the SBIP + IMP intervention condition (both early and late in intervention). The results indicate that using an IMP helps to establish an agreement between IPV offenders and facilitators on BIPs' objectives and tasks. Reaching this agreement is a challenge in BIPs, as IPV offenders tend to minimize or fail to recognize acts of violence for which they have been convicted (Flinck & Paavilainen, 2008; Lila, Oliver, Catalá-Miñana, & Conchell, 2014; Martín-Fernández et al., 2018; Murphy & Maiuro, 2009; Weber et al., 2019). For example, they do not recognize that their anger is problematic, and tend to attribute responsibility for their abusive actions to their partner or the legal system, believing that they have been treated unfairly and that they are the main aggrieved party (DiGiuseppe et al., 1994; Murphy & Maiuro, 2009; Vitoria-Estruch et al., 2017). In addition, most of these men are court referred or enter intervention as a result of external pressure (Daly & Pelowski, 2000; Velonis et al., 2016). Motivational strategies seem to overcome some of these difficulties by establishing a collaborative environment and helping participants to establish self-determined goals through the exploration of potential benefits of change (Lee et al., 2014; Miller & Rollnick, 2002). Regarding bond, Safran and Muran (2000) stressed the importance of participants' trust in facilitators' ability to help them throughout the intervention. Among the basic intervention principles of motivational strategies, such as the IMP, is the acceptance of participant resistance as part of the change process (Murphy & Maiuro, 2009). This attitude of accepting resistance and supporting participants' self-efficacy facilitates and strengthens the bond between a facilitator and a participant.

As for protherapeutic behaviors, our results showed that assumption of responsibility was significantly higher in the SBIP + IMP intervention condition (both early and late in intervention). Participants in the SBIP + IMP condition recognized more frequently within the group their responsibility for the violence, its consequences on people around them, and the need to make personal changes to avoid committing abusive acts in the future. Non-confrontational, non-judgmental, and empathetic listening qualities of motivational strategies may explain this increase in participants' assumption of responsibility (Musser et al., 2008; Taft & Murphy, 2007). In fact, the framework of motivational interviewing is that participants feel accepted despite the presence of some unacceptable behaviors (Murphy & Maiuro, 2009). Concerning early in intervention participant role behavior, offenders in the SBIP + IMP intervention condition showed significantly greater efforts to help other members of the group to change and to assume responsibility for their behavior. For example, participants in the SBIP + IMP group more frequently confronted other group members' comments about avoiding responsibility by blaming their partner or the legal system, or reinforced assumptions of responsibility of other group members (Gracia, 2014; Henning & Holdford, 2006; Lila, Oliver, Catalá-Miñana, Galiana, et al., 2014; Martín-Fernández et al., 2018). Regarding group value, participants in the SBIP + IMP intervention condition were

more likely to make positive assessments of the group, and the intervention in general, both early and late in intervention.

The results of our study highlight the importance of using motivational strategies in order to increase their impact on both participant-facilitator working alliance and participants' protherapeutic behaviors. Our study pulls ahead previous research on the importance of motivational strategies to increase the effectiveness of BIPs (Musser et al., 2008) at least in three aspects. First, in our study, motivational strategies were implemented throughout the intervention program. Other studies tend to implement motivational strategies only at pre-intervention stage. However, as our results showed, a more extensive implementation of motivational strategies can lead to more long-lasting gains (Crane & Eckhardt, 2013; Musser et al., 2008; Santirso et al., 2020; Stuart et al., 2013). This idea is also supported by research on other behavior change interventions. For example, a systematic review of the effectiveness of motivational interviewing on substance abuse, gambling, health-related behaviors, and engagement in intervention showed a positive association between the motivational intervention dose and its efficacy (Lundahl et al., 2010). Second, in our study we assess both protherapeutic behaviors and working alliance using a systematic observational methodology conducted by external observers. This methodological approach allows researchers to address important limitations that arise when using offenders' self-reports (i.e., impression management or social desirability). This approach also overcomes potential biases in facilitators' reports (Bennett, 2007; Gracia et al., 2015; Santirso et al., 2018). For example, previous research reports different results depending on the source of information, such as participants or facilitators (Musser et al., 2008). Third, facilitators used in our study were psychologists with at least two years' experience in managing BIPs. Facilitators' experience can be a major factor in the implementation quality of motivational strategies (Hamel et al., in press).

As for limitations, although manuals for each condition were used to ensure the content of and adherence to the protocol, and facilitators were regularly supervised, the IMP protocol was not quantitatively rated by an external observer. Also, this study was conducted with IPV offenders court-mandated to a community-based BIP. This could limit the generalizability of our results to other population samples such as men attending voluntary programs, men imprisoned for IPV, women who have committed IPV, or individuals with severe substance abuse problems or psychological disorders. Despite these limitations, this is the first RCT evaluating the effectiveness of a motivational strategy implemented throughout the intervention program—IMP—on the participant-facilitator working alliance and participants' protherapeutic behaviors. Our findings have important practical implications, as our results clearly showed that a motivational strategy tool such as the IMP improves key intervention processes (i.e., working alliance and protherapeutic behaviors) in BIPs, therefore increasing their effectiveness.

Conflict of Interest

The authors of this article declare no conflict of interest.

References

- Alexander, P. C., Morris, E., Tracy, A., & Frye, A. (2010). Stages of change and the group treatment of batterers: A randomized clinical trial. *Violence and Victims, 25*, 571-587. <https://doi.org/10.1891/0886-6708.25.5.571>
- Arce, R., Arias, E., Novo, M., & Fariña, F. (2020). Are interventions with batterer effectiveness? A meta-analytical review. *Psychosocial Intervention*. Advance online. <https://doi.org/10.5093/pi2020a11>
- Babcock, J., Armenti, N., Cannon, C., Lauve-Moon, K., Buttell, F., Ferreira, R., Cantos, A., Hamel, J., Kelly, D., Jordan, C., Lehmann, P., Leising, P. A., Murphy, C., O'Leary, K. D., Bannon, S., Salis, K. L., & Solano, I. (2016). Domestic violence perpetrator programs: A proposal for evidence-based standards in the United States. *Partner Abuse, 7*, 355-460. <https://doi.org/10.1891/1946-6560.7.4.355>
- Babcock, J. C., Green, C. E., & Robie, C. (2004). Does batterers' treatment work? A meta-analytic review of domestic violence treatment. *Clinical Psychology Review, 23*, 1023-1053. <https://doi.org/10.1016/j.cpr.2002.07.001>
- Bennett, L. W., Stoops, C., Call, C., & Flett, H. (2007). Program completion and re-arrest in a batterer intervention system. *Research on Social Work Practice, 17*, 42-54. <https://doi.org/10.1177/1049731506293729>
- Bordin, E. S. (1979). The generalizability of the psychoanalytic concept of the working alliance. *Psychotherapy: Theory, Research and Practice, 16*, 252-260. <https://doi.org/10.1037/h0085885>
- Brown, P. D., & O'Leary, K. D. (2000). Therapeutic alliance: Predicting continuance and success in group treatment for spouse abuse. *Journal of Consulting and Clinical Psychology, 68*, 340-345. <https://doi.org/10.1037/0022-006x.68.2.340>
- Cadsky, O., Hanson, R. K., Crawford, M., & Lalonde, C. (1996). Attrition from a male batterer treatment program: Client-treatment congruence and lifestyle instability. *Violence and Victims, 11*, 51-64. <https://doi.org/10.1891/0886-6708.11.1.51>
- Carbajosa, P., Catalá-Miñana, A., Lila, M., & Gracia, E. (2017). Differences in treatment adherence, program completion, and recidivism among batterer subtypes. *The European Journal of Psychology Applied to Legal Context, 9*, 93-101. <https://doi.org/10.1016/j.ejpal.2017.04.001>
- Carbajosa, P., Catalá-Miñana, A., Lila, M., Gracia, E., & Boira, S. (2017). Responsive versus treatment-resistant perpetrators in batterer intervention programs: Personal characteristics and stages of change. *Psychiatry, Psychology and Law, 24*, 936-950. <https://doi.org/10.1080/13218719.2017.1347933>
- Cheng, S. Y., Davis, M., Jonson-Reid, M., & Yaeger, L. (2019). Compared to what? A meta-analysis of batterer intervention studies using nontreated controls or comparisons. *Trauma, Violence, & Abuse, 1-16*. <https://doi.org/10.1177/1524838019865927>
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* (2nd ed). Erlbaum.
- Crane, C. A., & Eckhardt, C. I. (2013). Evaluation of a single-session brief motivational enhancement intervention for partner abusive men. *Journal of Counseling Psychology, 60*, 180-187. <https://doi.org/10.1037/a0032178>
- Crane, C. A., Schlauch, R. C., & Eckhardt, C. I. (2015). Dyadic violence and readiness to change among male intimate partner violence offenders. *Criminal Behaviour and Mental Health, 25*, 287-298. <https://doi.org/10.1002/cbm.1977>
- Daly, J. E., & Pelowski, S. (2000). Predictors of dropout among men who batter: A review of studies with implications for research and practice. *Violence and Victims, 15*, 137-160. <https://doi.org/10.1891/0886-6708.15.2.137>
- De Shazer, S., & Berg, I. K. (1997). 'What works?' Remarks on research aspects of solution-focused brief therapy. *Journal of Family Therapy, 19*, 121-124. <https://doi.org/10.1111/1467-6427.00043>
- DiGiuseppe, R., Tafrate, R., & Eckhardt, C. (1994). Critical issues in the treatment of anger. *Cognitive and Behavioral Practice, 1*, 111-132. [https://doi.org/10.1016/S1077-7229\(05\)80089-6](https://doi.org/10.1016/S1077-7229(05)80089-6)
- Eckhardt, C., Holtzworth-Munroe, A., Norlander, B., Sibley, A., & Cahill, M. (2008). Readiness to change, partner violence subtypes, and treatment outcomes among men in treatment for partner assault. *Violence and Victims, 23*, 446-475. <https://doi.org/10.1891/0886-6708.23.4.446>
- Eckhardt, C. I., Murphy, C. M., Whitaker, D. J., Sprunger, J., Dykstra, R., & Woodard, K. (2013). The effectiveness of intervention programs for perpetrators and victims of intimate partner violence. *Partner Abuse, 4*, 196-231. <https://doi.org/10.1891/1946-6560.4.2.196>
- Feder, L., & Wilson, D. B. (2005). A meta-analytic review of court-mandated batterer intervention programs: Can courts affect abusers' behavior? *Journal of Experimental Criminology, 1*, 239-262. <https://doi.org/10.1007/s11292-005-1179-0>
- Flinck, A., & Paavilainen, E. (2008). Violent behavior of men in their intimate relationships, as they experience it. *American Journal of Men's Health, 2*, 244-253. <https://doi.org/10.1177/1557988308321674>
- Gracia, E. (2014). Intimate partner violence against women and victim-blaming attitudes among Europeans. *Bulletin of the World Health Organization, 92*, 380-381. <https://doi.org/10.2471/BLT.13.131391>
- Gracia, E., Rodriguez, C. M., & Lila, M. (2015). Preliminary evaluation of an analog procedure to assess acceptability of intimate partner violence against women: The partner violence acceptability movie task. *Frontiers in Psychology, 6*, 1567. <https://doi.org/10.3389/fpsyg.2015.01567>
- Hamel, J., Cannon, C., Buttell, F., & Ferreira, R. (in press). A survey of IPV perpetrator treatment providers: Ready for evidence-based practice? *Partner Abuse*.
- Henning, K., & Holdford, R. (2006). Minimization, denial, and victim blaming by batterers: How much does the truth matter? *Criminal Justice and Behavior, 33*, 110-130. <https://doi.org/10.1177/0093854805282322>
- Jewell, L. M., & Wormith J. S. (2010). Variables associated with attrition from domestic violence treatment programs targeting male batterers: A meta-analysis. *Criminal Justice and Behavior, 37*, 1086-1113. <https://doi.org/10.1177/0093854810376815>

- Juarros-Basterrechea, J., Herrero, J., Fernández-Suárez, A., Pérez, B., & Rodríguez-Díaz, F. J. (2018). Are generalist batterers different from generally extra-family violent men? A study among imprisoned male violent offenders. *European Journal of Psychology Applied to Legal Context*, 10, 8-14. <https://doi.org/10.5093/ejpalc2018a1>
- Kistenmacher, B. R., & Weiss, R. L. (2008). Motivational interviewing as a mechanism for change in men who batter: A randomized controlled trial. *Violence and Victims*, 23, 558-570. <https://doi.org/10.1891/0886-6708.23.5.558>
- Langlands, R. L., Ward, T., & Gilchrist, E. (2009). Applying the Good Lives Model to male perpetrators of domestic violence. *Behaviour Change*, 26, 113-129. <https://doi.org/10.1375/bech.26.2.113>
- Lee, M. Y., Uken, A., & Sebold, J. (2014). Self-determined goals and treatment of domestic violence offenders: What if we leave it up to them? *Partner Abuse*, 5, 239-258. <http://doi.org/10.1891/1946-6560.5.3.239>
- Lila, M., Gracia, E., & Catalá-Miñana, A. (2018). Individualized motivational plans in batterer intervention programs: A randomized clinical trial. *Journal of Consulting and Clinical Psychology*, 86, 309-320. <http://doi.org/10.1037/ccp0000291>
- Lila, M., Martín-Fernández, M., Gracia, E., López-Ossorio, J. J., & González, J. L. (2019). Identifying key predictors of recidivism among offenders attending a batterer intervention program: A survival analysis. *Psychosocial Intervention*, 28, 157-167. <https://doi.org/10.5093/pi2019a19>
- Lila, M., Oliver, A., Catalá-Miñana, A., & Conchell, R. (2014). Recidivism risk reduction assessment in batterer intervention programs: A key indicator for program efficacy evaluation. *Psychosocial Intervention*, 23, 217-223. <https://doi.org/10.1016/j.psi.2014.11.001>
- Lila, M., Oliver, A., Catalá-Miñana, A., Galiana, L., & Gracia, E. (2014). The Intimate Partner Violence Responsibility Attribution Scale (IPVRAS). *European Journal of Psychology Applied to Legal Context*, 6, 29-36. <https://doi.org/10.5093/ejpalc2014a4>
- Lundahl, B. W., Kunz, C., Brownell, C., Tollefson, D., & Burke, B. L. (2010). A meta-analysis of motivational interviewing: Twenty-five years of empirical studies. *Research on Social Work Practice*, 20, 137-160. <https://doi.org/10.1177/1049731509347850>
- Martín-Fernández, M., Gracia, E., & Lila, M. (2018). Assessing victim-blaming attitudes in cases of intimate partner violence against women: Development and validation of the VB-IPVAW scale. *Psychosocial Intervention*, 27, 133-143. <https://doi.org/10.5093/pi2018a18>
- Martín-Fernández, M., Gracia, E., Marco, M., Vargas, V., Santirso, F. A., & Lila, M. (2018). Measuring acceptability of intimate partner violence against women: Development and validation of the A-IPVAW scale. *European Journal of Psychology Applied to Legal Context*, 10, 26-34. <https://doi.org/10.5093/ejpalc2018a3>
- Mbilinyi, L. F., Neighbors, C., Walker, D. D., Roffman, R. A., Zegree, J., Edleson, J., & O'Rourke, A. (2011). A telephone intervention for substance-using adult male perpetrators of intimate partner violence. *Research on Social Work Practice*, 21, 43-56. <https://doi.org/10.1177/1049731509359008>
- Miller, W. R., & Rollnick, S. (2002). *Motivational interviewing: Preparing people for change*. Guilford.
- Murphy, C. M., & Baxter, V. A. (1997). Motivating batterer to change in the treatment context. *Journal of Interpersonal Violence*, 12, 607-619. <https://doi.org/10.1177/088626097012004009>
- Murphy, C. M., Eckhardt, C. I., Clifford, J. M., Lamotte, A. D., & Meis, L. A. (2017). Individual versus group cognitive-behavioral therapy for partner-violent men: A preliminary randomized trial. *Journal of Interpersonal Violence*, 1-23. <https://doi.org/10.1177/0886260517705666>
- Murphy, C. M., & Maiuro, R. D. (2009). *Motivational interviewing and stages of change in intimate partner violence*. Springer Publishing Company.
- Musser, P. H., Semiatin, J. N., Taft, C. T., & Murphy, C. M. (2008). Motivational interviewing as a pregroup intervention for partner-violent men. *Violence and Victims*, 23, 539-557. <https://doi.org/10.1891/0886-6708.23.5.539>
- Olver, M. E., Stockdale, K. C., & Wormith, J. S. (2011). A meta-analysis of predictors of offender treatment attrition and its relationship to recidivism. *Journal of Consulting and Clinical Psychology*, 79, 6-21. <https://doi.org/10.1037/a0022200>
- Prochaska, J. O., & DiClemente, C. C. (1982). Transtheoretical therapy: Toward a more integrative model of change. *Psychotherapy: Theory, Research and Practice*, 19, 276-288. <https://doi.org/10.1037/h0088437>
- Prochaska, J. O., DiClemente, C. C., & Norcross, J. C. (1992). In search of how people change: Applications to addictive behaviors. *American Psychologist*, 47, 1102-1114. <https://doi.org/10.1037/0003-066x.47.9.1102>
- Romero-Martínez, Á., Lila, M., Gracia, E., & Moya-Albiol, L. (2019). Improving empathy with motivational strategies in batterer intervention programmes: Results of a randomized controlled trial. *British Journal of Clinical Psychology*, 58, 125-139. <https://doi.org/10.1111/bjc.12204>
- Rondeau, G., Brodeur, N., Brochu, S., & Lemire, G. (2001). Dropout and completion of treatment among spouse abusers. *Violence and Victims*, 16, 127-143. <https://doi.org/10.1891/0886-6708.16.2.127>
- Safran, J. D., & Muran, J. C. (2000). Resolving therapeutic alliance ruptures: Diversity and integration. *Journal of Clinical Psychology*, 56, 233-243. [https://doi.org/10.1002/\(SICI\)1097-4679\(200002\)56:2<233::AID-JCLP9>3.0.CO;2-3](https://doi.org/10.1002/(SICI)1097-4679(200002)56:2<233::AID-JCLP9>3.0.CO;2-3)
- Santirso, F. A., Gilchrist, G., Lila, M., & Gracia, E. (2020). Motivational strategies for intimate partner violence offender treatment: A systematic review and meta-analysis of randomized controlled trials. *Psychosocial Intervention*, 29. <https://doi.org/10.5093/pi2020a13>
- Santirso, F. A., Martín-Fernández, M., Lila, M., Gracia, E., & Terreros, E. (2018). Validation of the working alliance inventory-observer short version with male intimate partner violence offenders. *International Journal of Clinical and Health Psychology*, 18, 152-161. <https://doi.org/10.1016/j.ijchp.2018.02.003>
- Scott, K., King, C., McGinn, H., & Hosseini, N. (2011). Effects of motivational enhancement on immediate outcomes of batterer intervention. *Journal of Family Violence*, 26, 139-149. <https://doi.org/10.1007/s10896-010-9353-1>
- Semiatin, J. N., Murphy, C. M., & Elliott, J. D. (2013). Observed behavior during group treatment for partner-violent men: Acceptance of responsibility and promotion of change. *Psychology of Violence*, 3, 126-139. <https://doi.org/10.1037/a0029846>
- Smedslund, G., Dalsbø, T. K., Steiro, A., Winsvold, A., & Clench-Aas, J. (2011). Cognitive behavioural therapy for men who physically abuse their female partner. *Cochrane Database of Systematic Reviews*. <https://doi.org/10.1002/14651858.CD006048.pub2>
- Stuart, G. L., Shorey, R. C., Moore, T. M., Ramsey, S. E., Kahler, C. W., O'Farrell, T. J., Strong, D. R., Temple, J. R., & Monti, P. M. (2013). Randomized clinical trial examining the incremental efficacy of a 90-minute motivational alcohol intervention as an adjunct to standard batterer intervention for men. *Addiction*, 108, 1376-1384. <https://doi.org/10.1111/add.12142>
- Stuart, G. L., Temple, J., & Moore, T. (2007). Improving batterer intervention programs through theory-based research. *Journal of the American Medical Association*, 298, 560-562. <https://doi.org/10.1001/jama.298.5.560>
- Taft, C. T. & Murphy, C. M. (2007). The working alliance in intervention for partner violence perpetrators: Recent research and theory. *Journal of Family Violence*, 22, 11-18. <https://doi.org/10.1007/s10896-006-9053-z>
- Taft, C. T., Murphy, C. M., Elliott, J. D., & Morrel, T. M. (2001). Attendance-enhancing procedures in group counselling for domestic abusers. *Journal of Counselling Psychology*, 48, 151-160. <https://doi.org/10.1037/0022-0167.48.1.151>
- Taft, C. T., Murphy, C. M., King, D. W., Musser, P. H., & DeDeyn, J. M. (2003). Process and treatment adherence factors in group cognitive-behavioral therapy for partner violent men. *Journal of Consulting and Clinical Psychology*, 71, 812-820. <https://doi.org/10.1037/0022-006X.71.4.812>
- Taft, C. T., Murphy, C. M., Musser, P. H., & Remington, N. A. (2004). Personality, interpersonal, and motivational predictors of the working alliance in group cognitive-behavioral therapy for partner-violent men. *Journal of Consulting and Clinical Psychology*, 72, 349-354. <https://doi.org/10.1037/0022-006X.72.2.349>
- Tichenor, V., & Hill, C. E. (1989). A comparison of six measures of working alliance. *Psychotherapy: Theory, Research, Practice, Training*, 26, 195-199. <https://doi.org/10.1037/h0085419>
- Velonis, A. J., Cheff, R., Finn, D., Davloor, W., & O'Campo, P. (2016). Searching for the mechanisms of change: A protocol for a realist review of batterer treatment programmes. *BMJ Open*, 6, 1-9. <https://doi.org/10.1136/bmjopen-2015-010173>
- Vitoria-Estruch, S., Romero-Martínez, Á., Ruiz-Robledillo, N., Sariñana-González, P., Lila, M., & Moya-Albiol, L. (2017). The role of mental rigidity and alcohol consumption interaction on intimate partner violence: A Spanish study. *Journal of Aggression, Maltreatment & Trauma*, 26, 664-675. <https://doi.org/10.1080/10926771.2017.1300619>
- Ward, T. (2002). Good lives and the rehabilitation of offenders: Promises and problems. *Aggression and Violent Behavior*, 7, 513-528. [https://doi.org/10.1016/S1359-1789\(01\)00076-3](https://doi.org/10.1016/S1359-1789(01)00076-3)
- Weber, E. N., Taylor, A. R., Cantos, A. L., Amado, B. G., & O'Leary, K. D. (2019). Exploring typology categorizations of male perpetrators: A methodology study. *European Journal of Psychology Applied to Legal Context*, 11, 93-97. <https://doi.org/10.5093/ejpalc2019a5>
- Woodin, E. M., & O'Leary, K. D. (2010). A brief motivational intervention for physically aggressive dating couples. *Prevention Science*, 11, 371-383. <https://doi.org/10.1007/s11121-010-0176-3>
- Zalmanowitz, S. J., Babins-Wagner, R., Rodger, S., Corbett, B. A., & Leschied, A. (2013). The association of readiness to change and motivational interviewing with treatment outcomes in males involved in domestic violence group therapy. *Journal of Interpersonal Violence*, 28, 956-974. <https://doi.org/10.1177/0886260512459381>