



Parenting Stress Index-Short Form: Psychometric Properties of the Spanish Version in Mothers of Children Aged 0 to 8 Years

Gabriela R. Rivas, Ignacia Arruabarrena, and Joaquín de Paúl

University of the Basque Country, Donostia-San Sebastián, Spain

ARTICLE INFO

Received 31 January 2020
Accepted 17 April 2020
Available online 22 June 2020

Keywords:

Parenting stress
Spanish adaptation
Factor structure
Psychometric properties

ABSTRACT

The Parenting Stress Index-Short Form (PSI-SF) is one of the most commonly used measures of parenting stress both in clinical and research contexts. The PSI-SF is a 36-item, self-report measure with three subscales: Parental Distress (PD), Parent-Child Dysfunctional Interaction (PCDI), and Difficult Child (DC). The objective of this study was to analyse the factor structure and psychometric properties of the Spanish version of PSI-SF. Two different samples ($N = 309$) of mothers with children under 8 years old participated in the study. The first sample comprised 203 mothers with difficulties managing their children's behaviour. The second sample comprised 106 mothers from the general population. Factor structure of the PSI-SF, convergent validity, and differences between groups were analysed. The expected three-factor structure was confirmed for both samples. Findings suggested that the total PSI-SF scale and the three subscales had adequate internal consistency and convergent validity. Differences between both samples, and between age and economic subgroups in the first sample were tested. The Spanish version of the PSI-SF can be considered an adequate measure of parenting stress in mothers of children under 8 years old with difficulties to manage their children's behaviour. Further studies with extended samples from the general population are needed.

Propiedades psicométricas de la versión española del *Parenting Stress Index-Short Form* en madres con niños o niñas de 0 a 8 años

RESUMEN

El *Parenting Stress Index-Short Form* (PSI-SF) es uno de los instrumentos más utilizados para evaluar el estrés parental tanto en el contexto clínico como en el de investigación. El PSI-SF es un autoinforme de 36 ítems con tres subescalas: malestar parental (PD), interacción disfuncional padre/madre-hijo (PCDI) y niño difícil (DC). El objetivo del estudio fue analizar la estructura factorial y las propiedades psicométricas de la versión española del PSI-SF. Dos muestras diferentes ($N = 309$) de madres con hijos o hijas menores de 8 años participaron en el estudio: la primera muestra estuvo compuesta por 203 madres con dificultades para gestionar la conducta de sus hijos o hijas y la segunda por 106 madres de la población general. Se analizó la estructura factorial del PSI-SF, la validez convergente y la diferencia entre grupos de madres, confirmándose la estructura de tres factores para las dos muestras. Los resultados sugieren que la escala completa del PSI-SF y las 3 subescalas presentaban una adecuada consistencia interna y validez convergente. Se analizaron las diferencias en todas las puntuaciones del PSI-SF entre ambas muestras y entre subgrupos (edad y situación económica) de la primera muestra. La versión española del PSI-SF puede considerarse un instrumento adecuado para medir el estrés parental en madres con niños o niñas menores de 8 años con dificultades para manejar su comportamiento. Se necesitan estudios con muestras más representativas de la población general.

Palabras clave:

Estrés parental
Adaptación española
Estructura factorial
Propiedades psicométricas

Parenting stress is a complex process that can be subsumed within the general stress model of Lazarus and Folkman (1984) and understood as the aversive psychological reaction that occurs when caregivers feel overwhelmed and perceive that they lack the skills required to cope with their parental role (Abidin, 1995; Deater-Deckard, 1998). Parenting stress is conceptually distinct from other

life stressors that a parent might experience (e.g., negative life events, financial problems), although they are frequently related (Holly et al., 2019).

Parenting stress has been found to be associated with parenting processes across all developmental periods. Research suggests that parenting stress tends to show stability and to decrease over time as

Cite this article as: Rivas, G. R., Arruabarrena, I., & de Paúl, J. (2021). Parenting stress index-short form: Psychometric properties of the Spanish version in mothers of children aged 0 to 8 years. *Psychosocial Intervention*, 30(1), 27-34. <https://doi.org/10.5093/pi2020a14>

Financing: This research was supported by a grant from the Ministry of Economy and Competitiveness (MINECO) of Spain (PSI2013-46272-P) and by a grant from the University of the Basque Country (UPV-EHU) (PIF2016/188). Correspondence: gabriela.rivas@ehu.eus (G. R. Rivas).

ISSN: 1132-0559/© 2021 Colegio Oficial de la Psicología de Madrid. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

the child becomes older, particularly when its initial levels are not very high (Neece et al., 2012; Stone et al., 2016; Williford et al., 2007).

Research has also shown parenting stress as a normative process that can affect every parent. However, it may be more severe for parents of children with clinically significant emotional, behavioural, or health issues (Crnic et al., 2005; Deater-Deckard & Panneton, 2017; Holly et al., 2019), and particularly challenging for families where parenting demands converge with negative situational circumstances (e.g., low-income) or personal difficulties (e.g., parents mental health problems, children with subclinical behaviour problems; Barroso et al., 2018; Menon et al., 2020). Higher levels of parenting stress have been found to be associated to depression and psychological difficulties in parents (Schleider et al., 2015; Theule et al., 2010; Thomason et al., 2014), behaviour problems and self-regulation difficulties in children (Anthony et al., 2005; Mackler et al., 2015; Mäntymaa et al., 2012; Stone et al., 2016), and negative interactions between parents and children (Dubois-Comtois et al., 2013; Gerdes et al., 2007; Van Steijn et al., 2014). Finally, there is evidence that parents' and children's factors contribute to parenting stress in a complex transactional process. Both contribute to parenting stress and at the same time are affected by it, having consequences for the well-being of parents and children (Crnic & Rose, 2017).

Considering the negative effects of parenting stress, its reduction constitutes a common and relevant goal of preventive and rehabilitative parenting programmes (Chen & Chan, 2016; Reyno & McGrath, 2006; Van Steijn et al., 2014), so reliable and valid measures are necessary. One of the most commonly instruments used in both clinical and research contexts is the Parenting Stress Index (PSI; Abidin, 1983), a 120-item self-report measure. Given its length, an abbreviated 36-item version was developed – the Parenting Stress Index-Short Form (PSI-SF; Abidin, 1983) – consisting of three subscales of twelve items each: Parental Distress (PD), Parent-Child Dysfunctional Interaction (PCDI), and Difficult Child (DC). The Parental Distress (PD) subscale captures the level of distress resulting from personal factors such as depression or conflict with a partner and life restrictions due to a parent's perception of his or her child-rearing competence. The Parent-Child Dysfunctional Interaction (PCDI) subscale assesses the extent to which a parent feels that his/her child is not meeting expectations and that interactions with the child are not reinforcing. The Difficult Child (DC) subscale measures a parent's view of his/her child's temperament, defiance, non-compliance, and demandingness. A total score of parenting stress is calculated by summing scores from the three subscales.

The Parenting Stress Index-Short Form (PSI-SF) has been used to measure parenting stress in parents from clinical and high-risk populations (Barbot et al., 2014; Crum & Moreland, 2017; Mackler et al., 2015; Vallotton et al., 2016), and to measure treatment effectiveness (Battagliese et al., 2015; Reyno & McGrath, 2006). PSI-SF has been translated to and applied in different languages like Italian (Miragoli et al., 2018), Spanish (Pérez & Menéndez, 2014), Portuguese (Seabra-Santos et al., 2016), and Finnish (Mäntymaa et al., 2012).

Excluding the study carried out by Abidin (1995), twenty studies have been found in Web of Science and PsycInfo databases analysing the psychometric properties of PSI-SF. Sixteen of those studies examined factor structure, yielding mixed findings. Confirmatory factor analyses in two of these studies found that the original three-factor model offered an adequate fit (Çekiç & Hamamci, 2018; Lee et al., 2008). Three other studies considered the three-factor model as the best choice due to its clinical value although with suboptimal fit indices (Lee et al., 2016; Reitman et al., 2002; Touchèque et al., 2016). Another group of studies involving both exploratory and confirmatory factor analyses concluded that the three-factor model was appropriate but removed some items (Dardas & Ahmad, 2014; Deater-Deckard & Scarr, 1996; Kang et al., 2017; Luo et al., 2019) or proposed two-factor (Haskett et al., 2006) or five-factor (McKelvey et al., 2009; Whiteside-Mansell et al., 2007; Zaidman-Zait et al., 2011)

models. Despite statistical analyses (confirmatory or exploratory factor analyses) and the amount of dimensions/factors proposed for the PSI-SF, all the studies reported moderate to high correlations between them, supporting the theoretical assumption that the relationship between PSI-SF's dimensions can be considered oblique.

Four studies have analysed factor structure or psychometric properties of the Spanish version of PSI-SF. Two of them used a sample of Spanish middle-class, married couples with infants aged between ten and thirty-nine-month old. Their findings suggested two different models for fathers and for mothers: whereas the original three-factor solution fitted the data for fathers (Díaz-Herrero et al., 2011), a two-factor model – labelled Childrearing Stress (CS), and Personal Distress (PD) – was proposed for mothers (Díaz-Herrero et al., 2010). A third study that administered PSI-SF's two-factor model proposed by Díaz-Herrero et al. (2010) to 109 Spanish at-risk mothers with one child aged below 12 years obtained satisfactory internal consistency and adequate discriminant validity indexes (Pérez-Padilla et al., 2015). Finally, an exploratory factor analysis of a fourth study carried out with a Chilean sample of 336 dyads consisting mostly of young, single mothers and their infants (age: $M = 84.8$ days, $SD = 78$ days) found that the PSI-SF shared the three-factor structure of the original, but proposed the elimination of two items (from the Parent-Child Dysfunctional Interaction and the Difficult Child subscales; Aracena et al., 2016).

As it has been seen, studies about the factor structure of the Parenting Stress Index-Short Form (PSI-SF) have not yielded consistent results. Several hypotheses can be proposed. First, differences may be linked to sample characteristics. For example, studies conducted with particular populations as younger, less educated, and poorer mothers of younger children (Aracena et al., 2016) or parents of children with autism spectrum disorders diagnosis (Zaidman-Zait et al., 2011) can yield different findings related to PSI-SF dimensions. Second, despite the possible effect of sample characteristics, differences between studies may be also linked to statistical analyses (e.g., confirmatory vs. exploratory factor analyses) used to explore PSI-SF dimensions.

Finally, studies analysing PSI-SF's convergent validity have found significant relationships between PSI-SF's total score and measures of family conflict, exposure to violence, and other negative life events; between the Parental Distress subscale and measures of depression or parental anxiety; and between the Difficult Child subscale and measures of child behaviour problems (Aracena et al., 2016; Barroso et al., 2016; Haskett et al., 2006; Lee et al., 2016; McKelvey et al., 2009; Pérez-Padilla et al., 2015; Reitman et al., 2002; Whiteside-Mansell et al., 2007; Zaidman-Zait et al., 2011).

The aim of the present study was to analyse the psychometric properties of the Spanish version of the Parenting Stress Index-Short Form (PSI-SF) with two groups of mothers with children aged 0 to 8 years old: mothers with significant problems to cope with their children's behaviour and mothers from the general population. The factor structure of PSI-SF Spanish version along with data on internal consistency and convergent validity were analysed as well as differences between groups.

Method

Participants

Two different samples of mothers of children aged under 8 years old participated in the study ($N = 309$). The first sample (clinical sample) consisted of 203 mothers with significant problems to cope with their children's behaviour recruited from family support and treatment programmes provided from Child Protection Services of the region of Gipuzkoa (Spain). The second (community sample) was a convenience sample consisting of 106 mothers from the general population of Gipuzkoa who were recruited via six schools that agreed to participate.

Table 1. Sociodemographic Characteristics of Mothers and Children from the Clinical and Community Samples

	Clinical Sample (n = 203)		Community Sample (n = 106)		t	df
	M (SD)		M (SD)			
Mother's age	35.1 (8.1)		39.9 (4.1)		5.68***	270
Child's age	5.3 (2.3)		6.3 (1.2)		3.78***	308
	n	%	n	%	χ^2	ΦV
Child's age					37.09***	.35
0-3 years	53	26.0	0	0.0		
4-6 years	82	40.2	70	66.0		
7-8 years	68	33.3	36	34.0		
Child's gender					1.34	.07
Male	128	62.7	59	55.7		
Female	76	37.3	47	44.3		
Country of origin					51.97***	.44
Spain	88	52.7	100	94.3		
Latin America	64	38.3	3	2.8		
Other	15	9.0	3	2.8		
Maternal education					132.12***	.69
Elementary	70	41.7	4	3.8		
High school	73	43.2	12	11.3		
Higher education	25	14.9	90	84.9		
Maternal employment					46.82***	.42
Permanent job	59	35.5	78	73.6		
Temporary job	26	15.7	17	16.0		
Unemployed	81	48.8	11	10.4		
Economic difficulties					69.31***	.50
Yes	86	51.2	3	2.8		
No	82	48.8	103	97.2		
Family composition					65.07***	.49
Two parents	71	42.8	97	91.5		
Single-parent	33	19.9	3	2.8		
Separated/divorced	62	37.3	6	5.7		

Note. M = mean; SD = standard deviation; df = degrees of freedom; χ^2 = chi-square; Φ = phi; V = Cramer's V.

*** $p < .001$.

As can be seen in Table 1, there were sample differences in socio-demographic characteristics. Compared to the community sample, mothers and children from the clinical sample were younger and included higher percentages of single-parent or separated/divorced families, mothers from other countries, with lower educational levels, and economic problems.

Instruments

Parenting Stress Index/Short Form (PSI-SF; Abidin, 1995). PSI-SF is a 36-item, self-report measure of parenting stress. It includes three subscales: Parental Distress (PD; e.g., "I feel trapped by my responsibilities as a parent", "I feel lonely and without friends"), Parent-Child Dysfunctional Interaction (PCDI; e.g., "Sometimes I feel my child doesn't like me and doesn't want to be close to me", "When I do things for my child I get the feeling that my efforts are not appreciated"), and Difficult Child (DC; e.g., "My child makes more demands on me than most children", "My child gets upset easily over the smallest thing"). Each subscale consists of 12 items rated from 1 (*strongly disagree*) to 5 (*strongly agree*), with subscales scores ranging from 12 to 60. A Total score is calculated by summing the three subscales scores, ranging from 36 to 180. Scores of 90 or above may indicate a clinical level of stress. Abidin (1995) reported Cronbach's alpha coefficients of .91 for PSI-SF total score, and .87, .80, and .85 for PD, PCDI, and DC subscales, respectively. Psychometric data obtained in the present study are presented in the Results section.

Beck Depression Inventory-II (BDI-II; Beck et al., 1996). BDI-II is a 21-item, self-report measure of depressive symptomatology.

This measure is appropriate for both psychiatric and normative populations. Responses are given using a four-point scale from 0 to 3 (0 = *I do not feel like a failure*, 1 = *I have failed more than I should have*, 2 = *As I look back, I see a lot of failures*, and 3 = *I feel I am a total failure as a person*), with scores ranging from 0 to 63 and higher scores indicating higher levels of depressive symptomatology. BDI-II has been shown to have adequate reliability (between .92 and .93 for internal consistency and $r = .93$ for test-retest reliability) as well as adequate construct validity (Beck et al., 1996). In the present study, internal consistency indices were satisfactory for both groups of mothers (Cronbach's alphas of .87 for the clinical sample, and .77 for the community sample).

Brief Child Abuse Potential Inventory (B-CAPI; Ondersma et al., 2005). B-CAPI is a self-report screening questionnaire composed by 34 items extracted from the Child Abuse Potential Inventory (CAP; Milner, 1986). Twenty-five items composed the Abuse scale that measures the risk of a parent to physically abuse their children (e.g., "I am often upset and do not know why", "Sometimes I feel lonely"), and two Validity scales: a three-item Random Response scale and a six-item Lie scale. Responses are in a binary scale (agree-disagree), so scores range between 0 to a maximum of 25 in the Abuse scale. Ondersma et al. (2005) indicated good internal consistency for the Abuse scale (KR20 = .89). In the present study, KR20 for the Abuse scale was computed for the clinical sample (.83) and for the community sample (.83). The 34 items used in this study were pulled out from the Spanish version of the CAP Inventory (De Paúl et al., 1999).

Eyberg Child Behaviour Inventory (ECBI; Eyberg & Pincus, 1999). ECBI is a parent-rating scale covering 36 common child disruptive be-

behaviours with two subscales. The Intensity subscale measures the frequency of a child's problem behaviour (e.g., "Acts defiant when told to do something", "Refuses to go to bed on time") on a seven-point scale, ranging from 1 to 7 with a minimum score of 36 and a maximum of 252. The Problem subscale measures the extent to which a parent finds their child's behaviour troublesome, which is rated on a binary scale (0 = No, 1 = Yes) with a range score from 0 to 36. Eyberg and Pincus (1999) reported high internal consistency for both Intensity and Problem subscales ($\alpha = .95$ and $KR20 = .94$, respectively). In the present study, both Intensity and Problem subscales showed high internal consistency for clinical ($\alpha = .91$ and $KR20 = .88$) and community samples ($\alpha = .88$ and $KR20 = .89$).

Procedure

The research design was approved by the Ethics Committee of the University of the Basque Country UPV/EHU (Spain). All participant mothers were informed of study goals and gave informed consent. Mothers in the clinical sample were informed by Child Protection Services caseworkers and completed the instruments at baseline and 6-months later in the presence of a trained clinical psychologist. Mothers in the community sample were informed by their children's school directors, collected the instruments from the school, completed them at home, and returned in a sealed envelope.

Data Analysis

Analyses were performed using IBM SPSS Statistics Version 24.0, Mplus 7.11, and R Studio. Preliminary analyses were conducted to examine data characteristics. Multivariate normality was estimated by Mardia's multivariate skewness and kurtosis tests (Mardia, 1970).

Confirmatory factor analysis (CFA) was used to examine PSI-SF factor structure

using weighted least squares mean and variance adjusted (WLSMV) estimation method. This method is recommended in non-normally distributed data with severe floor or ceiling effects (Brown, 2015). When the WLSMV estimator is used, missing data are treated with pairwise deletion, which is acceptable when the amount of missing data is minimal (Kline, 2011). In the present study, less than 1% of responses per PSI-SF item in both samples were missing.

Multiple fit indices were examined: root mean square error of approximation (RMSEA) values below .08 represent acceptable fit, comparative fit index (CFI) and Tucker-Lewis index (TLI) values between .90 and .95 represent reasonable model fit, and values above .95 represent excellent model fit (Brown, 2015).

Internal consistency was examined by computing Cronbach's alpha, McDonald's omega, and omega hierarchical coefficients for the PSI-SF as a whole and for the three subscales. Cronbach's alpha is less reliable in multidimensional measures and require equal factor loadings (Viladrich et al., 2017). Therefore, omega coefficients were also calculated using R Studio software.

Convergent validity was assessed by computing Spearman correlations between the Parental Distress subscale of the PSI-SF, and both the Beck Depression Inventory-II (BDI-II) and the Brief Child Abuse Potential Inventory (B-CAPI) scores, and between the Difficult Child subscale of the PSI-SF and the Eyberg Child Behaviour Inventory (ECBI) scores.

Measurement invariance (MI) between both samples and between subgroups of the clinical sample, child age, gender, and economic difficulties was intended to test. However, it was not possible to calculate MI because groups did not contained the same number of categories per item. For informative purposes, MANOVAs were conducted to test PSI-SF differences between clinical and community samples, and between subgroups in the clinical sample.

Differences between baseline and 6-months PSI-SF scores were analysed in the clinical sample with a repeated measures MANOVA.

Results

Preliminary Analysis

Analyses of item distribution of PSI-SF indicated violations of univariate normality in some items for both samples (see Table 3). Additionally, Mardia's multivariate skewness and kurtosis test was statistically significant ($p < .01$), suggesting a violation of multinormality in both samples.

PSI-SF Factor Structure and Reliability

Clinical sample. Three structural models of the PSI-SF were examined using confirmatory factor analysis (CFA): the original three-factor model proposed by Abidin (1995), the two-factor model proposed by Diaz-Herrero et al. (2010), and a one-factor model using PSI-SF total score. Goodness-of-fit indices for the three models tested are summarized in Table 2.

Table 2. Confirmatory Factor Analysis (CFA) Model Fit Indexes for One, Two, and Three Factors of the PSI-SF for the Clinical and Community Samples

Sample	Factors	χ^2	df	RMSEA	90% CI	p	CFI	TLI
Clinical sample	One factor	1656.72***	594	.10	[.09, .10]	< .05	.84	.81
	Two factor	1284.01***	593	.07	[.07, .08]	< .05	.89	.88
	Three factor	1135.28***	591	.06	[.06, .08]	< .05	.91	.90
Community sample	One factor	1178.19***	594	.09	[.09, .11]	< .05	.80	.78
	Two factor	897.89***	593	.07	[.06, .08]	< .05	.90	.89
	Three factor	817.60***	591	.06	[.05, .07]	< .05	.92	.92

Note. χ^2 = chi-square goodness of fit statistic; df = degrees of freedom; RMSEA = root-mean-square error of approximation; CFI = comparative fit index; TLI = Tucker Lewis index.
*** $p < .001$.

The three-factor model provided the best fit to the data with acceptable goodness of fit indices (RMSEA = .07, CFI = .91, TLI = .90). CFA items loadings, standardized errors, factor correlations, and reliability coefficients are presented in Table 3.

Items loadings and standard errors presented adequate estimates with loadings higher than .30 and standard errors between .03 and .09. Although two items loadings were lower to .30 (items 22 and 31) they were significant to their factors, so no further analyses were made.

As can be observed in Table 3, correlations between PSI-SF three factors ranged between .53 and .70. Following Brown (2015), the three dimensions showed adequate discriminant validity and were not overlapping. Furthermore, all reliability coefficients were adequate for the three-factor model.

Community sample. CFA analysis conducted for the community sample obtained similar findings than those for the clinical sample (see Table 2). The three-factor model provided the best fit to the data (RMSEA = .06, CFI = .92, TLI = .92). Factor loadings of the three-factor model were all above .30 with adequate standard errors between .04 and .12. Adequate correlations between the three factors were obtained, with values ranging between .34 and .77. Moreover, reliability coefficients showed adequate values for the model (see Table 3).

PSI-SF Convergent Validity

Correlations between BDI-II (depressive symptomatology), B-CAPI (child abuse potential), and ECBI (child behaviour problems), and PSI-SF Total and subscales scores were analysed in both samples (see Table 4). Total PSI-SF and Parental Distress subscale scores were

Table 3. Confirmatory Factor Analysis (CFA) Standardized Factor Loadings, Descriptive Statistics for Each Item, and Reliability Coefficients of PSI-SF for the Clinical and Community Samples

	Clinical Sample						Community Sample						
	M	SD	S	K	Factor loadings	SE	M	SD	S	K	Factor loadings	SE	
Parental Distress (PD) scale													
Item 1	3.3	1.2	-0.4	-0.9	.575*	.057	2.3	1.0	0.7	-0.3	.660*	.064	
Item 2	3.1	1.3	-0.1	-1.3	.574*	.053	2.7	1.2	0.2	-1.3	.683*	.065	
Item 3	2.5	1.2	0.6	-0.7	.654*	.050	2.2	0.9	0.9	0.8	.660*	.056	
Item 4	2.3	1.2	0.9	-0.2	.740*	.039	1.8	0.9	1.4	2.4	.591*	.062	
Item 5	2.3	1.1	0.9	-0.0	.816*	.033	2.0	0.8	0.9	0.7	.625*	.061	
Item 6	2.1	1.1	1.1	0.4	.509*	.060	1.9	0.9	1.2	1.6	.481*	.068	
Item 7	2.8	1.2	0.1	-1.2	.622*	.050	1.7	0.8	1.3	3.2	.706*	.055	
Item 8	2.0	1.1	1.1	0.6	.551*	.064	2.2	1.2	0.7	-0.7	.468*	.059	
Item 9	2.0	1.0	1.2	1.0	.645*	.051	1.4	0.7	1.6	2.7	.753*	.058	
Item 10	1.8	0.9	1.1	1.0	.664*	.051	1.5	0.7	1.1	1.0	.797*	.055	
Item 11	2.7	1.2	0.2	-1.2	.542*	.050	2.2	1.1	0.8	-0.4	.820*	.044	
Item 12	2.4	1.2	0.6	-0.8	.631*	.048	1.8	0.9	1.3	1.5	.778*	.042	
Parent-Child Dysfunctional Interaction (PCDI) scale													
Item 13	1.7	0.8	1.6	3.1	.821*	.032	1.2	0.5	1.9	3.0	.811*	.058	
Item 14	1.5	0.7	1.9	5.3	.855*	.030	1.1	0.3	2.3	3.3	.881*	.054	
Item 15	1.7	0.9	1.6	3.2	.806*	.030	1.2	0.4	1.6	1.5	.835*	.058	
Item 16	1.9	1.0	1.1	0.5	.775*	.034	1.6	0.7	1.0	0.3	.689*	.064	
Item 17	1.6	0.8	1.4	2.0	.819*	.032	1.3	0.5	1.2	0.5	.918*	.042	
Item 18	2.0	1.1	1.0	0.1	.653*	.041	1.3	0.7	2.3	5.2	.812*	.059	
Item 19	1.7	0.9	1.4	1.6	.880*	.023	1.3	0.5	1.8	2.2	.858*	.047	
Item 20	1.8	1.0	1.5	1.9	.747*	.039	1.2	0.4	1.9	2.5	.850*	.057	
Item 21	2.2	1.1	0.9	0.0	.636*	.044	1.5	0.7	1.5	2.0	.600*	.077	
Item 22	3.0	1.1	-0.7	-0.5	.244*	.078	2.5	0.9	-0.7	-0.6	.360*	.095	
Item 23	1.8	1.0	1.4	1.1	.797*	.033	1.4	0.6	2.0	4.6	.776*	.061	
Item 24	2.5	1.2	0.3	-1.3	.602*	.050	1.6	0.7	0.9	0.6	.635*	.069	
Difficult Child (DC) scale													
Item 25	2.2	1.2	0.9	-0.0	.743*	.044	1.5	0.7	1.5	2.1	.684*	.066	
Item 26	2.1	1.0	0.9	-0.1	.637*	.052	1.6	0.7	1.3	1.5	.525*	.073	
Item 27	3.1	1.2	-0.1	-1.1	.620*	.047	1.8	0.9	1.0	0.5	.740*	.057	
Item 28	3.4	1.1	-0.8	-0.4	.492*	.053	3.1	1.2	-0.7	-1.1	.561*	.069	
Item 29	3.3	1.2	-0.4	-1.0	.556*	.051	2.3	1.2	0.5	-0.8	.646*	.057	
Item 30	2.8	1.2	0.1	-1.2	.721*	.039	2.1	1.0	0.5	-0.8	.667*	.065	
Item 31	2.5	1.3	0.6	-0.9	.252*	.073	2.1	1.2	0.9	-0.4	.465*	.084	
Item 32	3.5	1.1	-0.6	-0.4	.397*	.066	3.2	0.9	-0.5	0.3	.657*	.074	
Item 33	2.0	1.1	1.1	0.4	.469*	.067	1.4	0.6	1.5	1.2	.473*	.119	
Item 34	3.3	1.2	-0.4	-1.0	.589*	.048	2.8	1.2	0.1	-1.2	.422*	.080	
Item 35	2.0	1.1	1.0	0.1	.836*	.036	1.2	0.5	2.8	10	.834*	.078	
Item 36	2.2	1.2	0.9	-0.1	.759*	.042	1.5	0.7	1.3	1.4	.877*	.045	
			PD	PCDI					PD	PCDI			
			PCDI		.606*				PCDI		.591*		
			DC		.538*	.701*			DC		.338*	.768*	
			α	CI	ω	CI	ω_h	α	CI	ω	CI	ω_h	
			PD	.86	[.84, .89]	.87	[.84, .89]	.85	.85	[.81, .90]	.86	[.82, .90]	.85
			PCDI	.91	[.89, .93]	.91	[.90, .93]	.91	.86	[.82, .90]	.86	[.82, .90]	.84
			DC	.85	[.82, .88]	.85	[.82, .88]	.85	.79	[.74, .85]	.79	[.73, .85]	.79
			PSI-SF total	.93	[.92, .94]	.93	[.91, .94]	.89	.88	[.85, .91]	.88	[.85, .91]	.87

Note. S = skewness; K = kurtosis; SE = standardized errors; α = alpha; ω = omega; ω_h = hierarchical omega; CI = confidence interval 95%.

* $p < .05$.

strongly, positively correlated with BDI-II and B-CAPI scores in both samples, indicating that mothers reporting more parental distress also reported more depressive symptomatology and a higher risk for physical child abuse. Total PSI-SF and Difficult Child subscale scores were also strongly positively correlated with ECBI scores in both samples, indicating that mothers who reported greater stress due to having a difficult child also reported more child behaviour problems. In addition, Parent-Child Dysfunctional Interaction subscale scores, although with less strength, were also positively correlated with both BDI-II and ECBI scores in both samples.

PSI-SF Differences between Baseline and 6-Month Measures

The repeated-measures MANOVA (see Table 5) revealed statistically significant differences between baseline and 6-month scores for the total PSI-SF and the Parental Distress, Parent-Child Dysfunctional Interaction, and Difficult Child subscales in the clinical sample of mothers (Wilks' lambda = .68, $F(3, 138) = 21.65$, $p < .0001$). As expected, 6-month scores (after receiving support or treatment services) were lower than baseline scores.

Table 4. Spearman Correlations between PSI-SF Total and Subscales Scores, and Mother Depressive Symptomatology (BDI-II), Child Abuse Potential (B-CAPI) and Child Behaviour Problems (ECBI) Scores

	Clinical Sample				Community Sample			
	PSI-SF	PD	PCDI	DC	PSI-SF	PD	PCDI	DC
Mother depressive symptoms (BDI-II)	.511***	.617***	.342***	.295***	.531***	.673***	.366***	.194*
Child abuse potential (B-CAPI)	.456***	.581***	.304***	.223***	.406***	.503***	.248*	.159
Child behaviour- Intensity (ECBI)	.500***	.225***	.353***	.644***	.532***	.234*	.466***	.606***
Child behaviour-Problem (ECBI)	.538***	.305***	.345***	.629***	.492***	.395***	.377***	.425***

Note. PSI-SF = Parental Stress Index-Short Form Total; PD = Parental Distress subscale; PCDI = Parent-Child Dysfunctional Interaction subscale; DC = Difficult Child subscale.
* $p < .05$, *** $p < .001$.

Table 5. Comparisons of PSI-SF Total and Subscales Scores between (1) Clinical and Community Samples, (2) Baseline and 6-month Measures in the Clinical Sample, (3) Children's age in the Clinical Sample, and (4) Economic Difficulties in the Family in the Clinical Sample

		PSI-SF	PD	PCDI	DC
		<i>M (SD)</i>	<i>M (SD)</i>	<i>M (SD)</i>	<i>M (SD)</i>
(1) Samples	Clinical sample	85.42 (19.3)	29.66 (8.4)	23.69 (7.3)	32.07 (8.3)
	Community sample	66.45 (14.2)	23.89 (6.8)	17.52 (4.6)	25.5 (6.7)
	<i>F</i> (1, 308)	79.8658***	37.153***	61.479***	57.098***
	<i>d</i>	1.07	0.73	0.94	0.91
Clinical sample					
(2) Baseline and 6-month measures	Baseline	86.47 (20.0)	29.49 (8.4)	24.26 (7.5)	32.72 (8.3)
	6 month-	76.52 (18.8)	26.35 (7.9)	22.23 (7.0)	27.94 (7.3)
	<i>F</i> (1, 140)	49.601***	28.516***	13.159***	61.785***
	<i>d</i>	0.41	0.39	0.28	0.61
(3) Children's age	0-3 years	78.71 (19.7)	29.59 (9.6)	21.31 (6.7)	27.80 (7.8)
	4-6 years	88.37 (18.8)	29.59 (8.1)	25.22 (7.7)	33.56 (7.7)
	7-8 years	86.68 (18.5)	29.81 (8.0)	23.46 (6.8)	33.4 (8.1)
	<i>F</i> (1, 201)	4.691***	0.030	4.610***	10.601***
	η^2	.046	.000	.044	.095
	Bonferroni post-hoc	0-3 < 4-6 0-3 < 7-8 4-6 = 7-8	0-3 = 4-6 0-3 = 7-8 4-6 = 7-9	0-3 < 4-6 0-3 = 7-8 4-6 = 7-10	0-3 < 4-6 0-3 = 7-8 4-6 = 7-8
(4) Economic difficulties	Yes	84.06 (20.8)	30.81 (9.1)	23.02 (7.8)	30.22 (8.6)
	No	84.63 (18.3)	27.74 (7.1)	23.99 (7.3)	31.53 (8.2)
	<i>F</i> (1, 167)	0.036	5.827***	0.678	4.537***
	<i>d</i>	0.03	0.37	0.13	0.33

Note. PSI-SF = Parental Stress Index-Short Form Total; PD = Parental Distress subscale; PCDI = Parent-child Dysfunctional Interaction subscale; DC = Difficult Child subscale;
d = Cohen's *d* effect size.
*** $p < .001$.

Comparison between Clinical and Community samples

Statistically significant differences between the PSI-SF scores of both samples were found (Table 5). Mothers of the clinical sample obtained higher scores on the total PSI-SF, Parental Distress subscale, Parent-Child Dysfunctional Interaction subscale, and Difficult Child subscale scores (Wilks' lambda = .79, $F(3, 306) = 26.99$, $p < .0001$) than mothers from the community sample, suggesting that mothers with significant difficulties managing their children's behaviour felt more stress associated to their parenting role than mothers from the general population.

PSI-SF and Sociodemographic Variables

Differences in the PSI-SF scores based on sociodemographic characteristics (see Table 1) were assessed in both samples using two MANOVAs. Significant differences for child's age (Wilks' lambda = .87, $F(3, 398) = 4.64$, $p < .0001$) and economic difficulties (Wilks' lambda = .89, $F(3, 164) = 6.45$, $p < .0001$) were observed only in the clinical sample.

Differences between mothers with children of different ages were statistically significant for the total PSI-SF, Parent-Child Dysfunctional Interaction subscale, and Difficult Child subscale scores, but not for the Parental Distress subscale score (Table 5). Mothers of children

between 0-3 years old reported lower scores than mothers of children between 4-6 and 7-8 years old. No differences between mothers with children 4-6 years old and mothers with children 7-8 years old were observed.

Mothers who reported economic difficulties also reported significantly higher scores on Parental Distress and Difficult Child subscales than mothers no reporting economic difficulties. No statistically significant differences were observed between both groups on Parent-Child Dysfunctional Interaction subscale and Total PSI-SF scores.

Discussion

The purpose of the present study was to analyse factor structure and psychometric properties of the Spanish version of the Parenting Stress Index-Short Form (PSI-SF) with two different samples of mothers with children under 8 years old.

The results showed that the original PSI-SF three-factor model was the most appropriate for mothers with significant difficulties managing their children's behaviour as well as for mothers from the general population. Adequate internal consistency was found for the PSI-SF total score, and for the Parental Distress (PD), Parent-Child Dysfunctional Interaction (PCDI), and Difficult Child (DC) subscales. Further, correlations between the three subscales showed significant

values but lesser than .80, supporting their discriminant validity (Brown, 2015). Convergent validity with measures of depressive symptomatology, child abuse potential, and child behaviour problems also supported the PSI-SF three-factor model for both samples. In line with previous studies, the Parental Distress (PD) subscale was highly associated to mothers' depressive symptomatology, whereas the Difficult Child (DC) subscale showed a stronger association with mothers' reports of child behaviour problems (Barroso et al., 2016; Lee et al., 2016; Reitman et al., 2002).

Findings of our clinical sample were consistent with previous studies conducted with mothers of children aged between 0 to 12 years old of similar sociodemographic characteristics (low socioeconomic status), that concluded that PSI-SF three-factor model was the most adequate (Aracena et al., 2016; Lee et al., 2016; Reitman et al., 2002). Conversely, findings of our sample of mothers from the general population differed from those obtained by Díaz-Herrero et al. (2010), which supported a two-factor model. Differences between both studies can be related to sample characteristics—whereas the study by Díaz-Herrero et al. (2010) was conducted with mothers of children under 3 years old, the present study was conducted with mothers of children between 4 and 8 years old. Both used convenience samples, so additional studies with broader and representative samples are necessary for a better understanding of the psychometric characteristics of the PSI-SF Spanish version with mothers from the general population.

Taken together, our findings suggested that the Spanish version of the PSI-SF with three factors is appropriate to measure parenting stress in mothers having difficulties to manage their children's behaviour, and is also useful to detect changes following interventions designed to improve parenting skills. Also, the assessment of the three dimensions of parenting stress could be used to focus treatment strategies and for clinical decision-making.

The present study has some limitations that should be taken into account. First, small samples sizes did not allow testing invariance across samples and subgroups. Larger samples of mothers are necessary in order to confirm the influence of children's age and economic difficulties on parenting stress. Second, we only collected data from mothers, and differences in parenting stress between mothers and fathers can exist. The two studies that explored this issue yielded mixed results: whereas Deater-Deckard and Scarr (1996) did not find significant differences in PSI-SF scores, Delvecchio et al. (2015) found mothers reporting higher levels of parenting stress than fathers. These studies, however, did not test for measurement invariance (MI) across groups, a recommended analysis to test that the factor structure of an instrument is equivalent across groups and that is not conditioned by sample characteristics (for more information, see Putnick & Bornstein, 2016). Thus, further studies confirming the equivalence of PSI-SF factor structure across mothers and fathers are needed. This equivalence is required before conducting comparison analyses between groups. Only one study (Luo et al., 2019) analysed the factor structure of the PSI-SF and examined the measurement invariance (MI) across mothers and fathers in a community sample from China. They found that fathers reported significantly higher scores than mothers only in the Parent-Child Dysfunctional Interaction subscale (PCDI). However, the PSI-SF version of Luo et al. (2019) only included 15 items, so their findings cannot be generalized to the 36-item original version of the PSI-SF.

Based on our findings and the findings of previous studies with Spanish samples, it can be concluded that the Spanish version of the Parenting Stress Index-Short Form (PSI-SF) is useful to measure parenting stress with mothers with children under 8 years old. Further analyses with larger samples and including mothers and fathers are necessary to compare PSI-SF validity for the general population.

Conflict of Interest

The authors of this article declare no conflict of interest.

References

- Abidin, R. R. (1983). *The Parenting Stress Index*. Pediatric Psychology Press.
- Abidin, R. R. (1995). *Parenting Stress Index: Professional manual* (3rd ed.). Psychological Assessment Resources, Inc.
- Anthony, L. G., Anthony, B. J., Glanville, D. N., Naiman, D. Q., Waanders, C., & Shaffer, S. (2005). The relationships between parenting stress, parenting behaviour and preschoolers' social competence and behaviour problems in the classroom. *Infant and Child Development*, 14(2), 133-154. <https://doi.org/10.1002/icd.385>
- Aracena, M., Gómez, E., Undurraga, C., Leiva, L., Marinkovic, K., & Molina, Y. (2016). Validity and reliability of the Parenting Stress Index-Short Form (PSI-SF) applied to a Chilean sample. *Journal of Child and Family Studies*, 25(12), 3554-3564. <https://doi.org/10.1007/s10826-016-0520-8>
- Barbot, B., Crossman, E., Hunter, S. R., Grigorenko, E. L., & Luthar, S. S. (2014). Reciprocal influences between maternal parenting and child adjustment in a high-risk population: A 5-year cross-lagged analysis of bidirectional effects. *American Journal of Orthopsychiatry*, 84(5), 567-580. <https://doi.org/10.1037/ort0000012>
- Barroso, N. E., Hungerford, G. M., Garcia, D., Graziano, P. A., & Bagner, D. M. (2016). Psychometric properties of the Parenting Stress Index-Short Form (PSI-SF) in a high-risk sample of mothers and their infants. *Psychological Assessment*, 28(10), 1331-1335. <https://doi.org/10.1037/pas0000257>
- Barroso, N. E., Mendez, L., Graziano, P. A., & Bagner, D. M. (2018). Parenting stress through the lens of different clinical groups: A systematic review & meta-analysis. *Journal of Abnormal Child Psychology*, 46(3), 449-461. <https://doi.org/10.1007/s10802-017-0313-6>
- Battagliese, G., Caccetta, M., Luppino, O. I., Baglioni, C., Cardi, V., Mancini, F., & Buonanno, C. (2015). Cognitive-behavioral therapy for externalizing disorders: A meta-analysis of treatment effectiveness. *Behaviour Research and Therapy*, 75, 60-71. <https://doi.org/10.1016/j.brat.2015.10.008>
- Beck, A. T., Steer, R. A., & Brown, G. K. (1996). *Manual for the Beck Depression Inventory- II*. Psychological Corporation.
- Brown, T. A. (2015). *Confirmatory factor analysis for applied research* (2nd ed.). Guilford Press.
- Çekic, A., & Hamamci, Z. (2018). Adaptation of the Parenting Stress Index-Short Form into Turkish: A study of validity and reliability. *Anatolian Journal of Psychiatry*, 19(1), 63-70. <https://doi.org/10.5455/apd.263093>
- Chen, M., & Chan, K. L. (2016). Effects of parenting programs on child maltreatment prevention: A meta-analysis. *Trauma, Violence, and Abuse*, 17(1), 88-104. <https://doi.org/10.1177/1524838014566718>
- Crníc, K., Gaze, C., & Hoffman, C. (2005). Cumulative parenting stress across the preschool period: Relation to maternal parenting and child behaviour at age 5. *Infant and Child Development*, 14, 117-132. <https://doi.org/10.1002/icd>
- Crníc, K., & Rose, E. (2017). Parenting stress and parental efficacy. In K. Deater-Deckard & R. Panneton (Eds.), *Parental stress and early child development: Adaptive and maladaptive outcomes* (pp. 263-284). Springer International Publishing.
- Crum, K. I., & Moreland, A. D. (2017). Parental stress and children's social and behavioral outcomes: The role of abuse potential over time. *Journal of Child and Family Studies*, 26(11), 3067-3078. <https://doi.org/10.1007/s10826-017-0822-5>
- Dardas, L. A., & Ahmad, M. M. (2014). Psychometric properties of the Parenting Stress Index with parents of children with autistic disorder. *Journal of Intellectual Disability Research*, 58(6), 560-571. <https://doi.org/10.1111/jir.12053>
- De Paúl, J., Arruabarrena, I., Múgica, P., & Milner, J. (1999). Validation of the Spanish version of the Child Abuse Potential Inventory. *Estudios de Psicología*, 20(63-64), 55-72. <https://doi.org/10.1174/02109399960256766>
- Deater-Deckard, K. (1998). Parenting stress and child adjustment: Some old hypotheses and new questions. *Clinical Psychology: Science and Practice*, 5(3), 314-332. <https://doi.org/10.1111/j.1468-2850.1998.tb00152.x>
- Deater-Deckard, K., & Panneton, R. (2017). Unearthing the developmental and intergenerational dynamics of stress in parent and child functioning. In K. Deater-Deckard & R. Panneton (Eds.), *Parental stress and early child development: Adaptive and maladaptive outcomes* (pp. 1-11). Springer International Publishing.
- Deater-Deckard, K., & Scarr, S. (1996). Parenting stress among dual-earner mothers and fathers: Are there gender differences? *Journal of Family Psychology*, 10(1), 45-59. <https://doi.org/10.1037/0893-3200.10.1.45>
- Delvecchio, E., Sciandra, A., Finos, L., Mazzeschi, C., & Di Riso, D. (2015). The role of co-parenting alliance as a mediator between trait anxiety, family system maladjustment, and parenting stress in a sample of non-clinical Italian parents. *Frontiers in Psychology*, 6, 1-8. <https://doi.org/10.3389/fpsyg.2015.01177>

- Díaz-Herrero, Á., de la Nuez, A. G. B., Pina, J. A. L., Pérez-López, J., & Martínez-Fuentes, M. T. (2010). Estructura factorial y consistencia interna de la versión española del Parenting Stress Index-Short Form. *Psicothema*, 22(4), 1033-1038.
- Díaz-Herrero, Á., López-Pina, J. A., Pérez-López, J., de la Nuez, A. G. B., & Martínez-Fuentes, M. T. (2011). Validity of the Parenting Stress Index-Short Form in a sample of Spanish fathers. *The Spanish Journal of Psychology*, 14(2), 990-997. https://doi.org/10.5209/rev_SJOP.2011.v14.n2.44
- Dubois-Comtois, K., Moss, E., Cyr, C., & Pascuzzo, K. (2013). Behavior problems in middle childhood: The predictive role of maternal distress, child attachment, and mother-child interactions. *Journal of Abnormal Child Psychology*, 41(8), 1311-1324. <https://doi.org/10.1007/s10802-013-9764-6>
- Eyberg, S. M., & Pincus, D. (1999). *Eyberg Child Behavior Inventory and Sutter-Eyberg Student Behavior Inventory-Revised: Professional manual*. Psychological Assessment Resources, Inc.
- Gerdes, A. C., Hoza, B., Arnold, L. E., Pelham, W. E., Swanson, J. M., Wigal, T., & Jensen, P. S. (2007). Maternal depressive symptomatology and parenting behavior: Exploration of possible mediators. *Journal of Abnormal Child Psychology*, 35(5), 705-714. <https://doi.org/10.1007/s10802-007-9134-3>
- Haskett, M. E., Ahern, L. S., Ward, C. S., & Allaire, J. C. (2006). Factor structure and validity of the Parenting Stress Index-Short Form. *Journal of Clinical Child & Adolescent Psychology*, 35(2), 302-312. https://doi.org/10.1207/s15374424jccp3502_14
- Holly, L. E., Fenley, A. R., Kritikos, T. K., Merson, R. A., Abidin, R. R., & Langer, D. A. (2019). Evidence-based update for parenting stress measures in clinical samples. *Journal of Clinical Child and Adolescent Psychology*, 48(5), 685-705. <https://doi.org/10.1080/15374416.2019.1639515>
- Kang, S.-K., Choi, H.-J., & Mi-Ra, C. (2017). A study of validity on the Parenting Stress Index-Short Form (PSI-SF) for grandmothers of infants and toddlers. *The Journal of Future Early Childhood Education*, 24(2), 101-125. <https://doi.org/10.22155/JFECE.24.2.101.125>
- Kline, R. B. (2011). *Principles and practice of structural equation modeling*. Guilford Press.
- Lazarus, R., & Folkman, S. (1984). *Stress, appraisal and coping*. Springer.
- Lee, K. S., Chung, K. M., Park, J. A., & Kim, H. J. (2008). Reliability and validity study for the Korean version of Parenting Stress Index-Short Form (K-PSI-SF). *The Korean Journal of Woman Psychology*, 13(3), 363-377. <https://doi.org/10.18205/kpa.2008.13.3.007>
- Lee, S. J., Gopalan, G., & Harrington, D. (2016). Validation of the Parenting Stress Index-Short Form with minority caregivers. *Research on Social Work Practice*, 26(4), 429-440. <https://doi.org/10.1177/1049731514554854>
- Leitão, S., Pimentel, M., Almeida, M., & Moura-Ramos, M. (2016). Incredible Years parent training: What changes, for whom, how, for how long? *Journal of Applied Developmental Psychology*, 44, 93-104. <https://doi.org/10.1016/j.appdev.2016.04.004>
- Luo, J., Wang, M.-C., Gao, Y., Zeng, H., Yang, W., Chen, W., Shouying, Z. S., & Qi, S. (2019). Refining the Parenting Stress Index-Short Form (PSI-SF) in Chinese parents. *Assessment*, 81(1), 1-16. <https://doi.org/10.1177/1073191119847757>
- Mackler, J. S., Kelleher, R. T., Shanahan, L., Calkins, S. D., Keane, S. P., & O'Brien, M. (2015). Parenting stress, parental reactions, and externalizing behavior from ages 4 to 10. *Journal of Marriage and Family*, 77(2), 388-406. <https://doi.org/10.1111/jomf.12163>
- Mäntymaa, M., Puura, K., Luoma, I., Latva, R., Salmelin, R. K., & Tamminen, T. (2012). Predicting internalizing and externalizing problems at five years by child and parental factors in infancy and toddlerhood. *Child Psychiatry & Human Development*, 43(2), 153-170. <https://doi.org/10.1007/s10578-011-0255-0>
- Mardia, K. (1970). Measures of multivariate skewness and kurtosis with applications. *Biometrika*, 57, 519-530.
- McKelvey, L. M., Whiteside-Mansell, L., Faldowski, R. A., Shears, J., Ayoub, C., & Hart, A. D. (2009). Validity of the Short Form of the Parenting Stress Index for fathers of toddlers. *Journal of Child and Family Studies*, 18(1), 102-111. <https://doi.org/10.1007/s10826-008-9211-4>
- Menon, M., Fauth, R. C., & Easterbrooks, M. A. (2020). Exploring trajectories of young mothers' parenting stress in early childhood: Associations with protective factors and psychological vulnerabilities. *Parenting*, 00(0), 1-29. <https://doi.org/10.1080/15295192.2020.1715683>
- Milner, J. S. (1986). *The Child Abuse Potential Inventory: Manual* (2nd ed.). Psytec.
- Miragoli, S., Balzarotti, S., Camisasca, E., & Di Blasio, P. (2018). Parents' perception of child behavior, parenting stress, and child abuse potential: Individual and partner influences. *Child Abuse & Neglect*, 84(1), 146-156. <https://doi.org/10.1016/j.chiabu.2018.07.034>
- Neece, C. L., Green, S. A., & Baker, B. L. (2012). Parenting stress and child behavior problems: A transactional relationship across time. *American Journal on Intellectual and Developmental Disabilities*, 117(1), 48-66. <https://doi.org/10.1352/1944-7558-117.1.48>
- Ondersma, S. J., Chaffin, M., Simpson, S., & LeBreton, J. (2005). The Brief Child Abuse Potential Inventory: Development and validation. *Journal of Clinical Child and Adolescent Psychology*, 34, 301-311.
- Pérez, J., & Menéndez, S. (2014). Un análisis tipológico del estrés parental en familias en riesgo psicosocial. *Salud Mental*, 37(1), 27-34.
- Pérez-Padilla, J., Menéndez, S., & Lozano, O. (2015). Validity of the Parenting Stress Index-Short Form in a Sample of At-Risk Mothers. *Evaluation Review*, 39(4), 428-446. <https://doi.org/10.1177/0193841X15600859>
- Putnick, D. L., & Bornstein, M. H. (2016). Measurement invariance conventions and reporting: The state of the art and future directions for psychological research. *Developmental Review*, 41, 71-90. <https://doi.org/10.1016/j.dr.2016.06.004>
- Reitman, D., Currier, R. O., & Stickle, T. R. (2002). A critical evaluation of the Parenting Stress Index-Short Form (PSI-SF) in a head start population. *Journal of Clinical Child & Adolescent Psychology*, 31(3), 384-392. https://doi.org/10.1207/S15374424JCCP3103_10
- Reyno, S. M., & McGrath, P. J. (2006). Predictors of parent training efficacy for child externalizing behavior problems – a meta-analytic review. *Journal of Child Psychology and Psychiatry*, 47(1), 99-111. <https://doi.org/10.1111/j.1469-7610.2005.01544.x>
- Schleider, J. L., Patel, A., Krumholz, L., Chorpita, B. F., & Weisz, J. R. (2015). Relation between parent symptomatology and youth problems: multiple mediation through family income and parent-youth stress. *Child Psychiatry and Human Development*, 46(1), 1-9. <https://doi.org/10.1007/s10578-014-0446-6>
- Seabra-Santos, M. J., Gaspar, M. F., Azevedo, A. F., Homem, T. C., Guerra, J., Martins, V., Leitao, S., Pimentel, M., Almeida, M., & Moura-Ramos, M. (2016). Incredible Years parent training: What changes, for whom, how, for how long? *Journal of Applied Developmental Psychology*, 44, 93-104. <https://doi.org/10.1016/j.appdev.2016.04.004>
- Stone, L. L., Mares, S. H. W., Otten, R., Engels, R. C. M. E., & Janssens, J. M. A. M. (2016). The co-development of parenting stress and childhood internalizing and externalizing problems. *Journal of Psychopathology and Behavioral Assessment*, 38(1), 76-86. <https://doi.org/10.1007/s10862-015-9500-3>
- Theule, J., Wiener, J., Tannock, R., & Jenkins, J. M. (2010). Parenting stress in families of children with ADHD. A meta-analysis. *Journal of Emotional and Behavioral Disorders*, 20(1), 1-15. <https://doi.org/10.1177/1063426610387433>
- Thomason, E., Volling, B. L., Flynn, H. A., McDonough, S. C., Marcus, S. M., Lopez, J. F., & Vazquez, D. M. (2014). Parenting stress and depressive symptoms in postpartum mothers: Bidirectional or unidirectional effects? *Infant Behavior and Development*, 37(3), 406-415. <https://doi.org/10.1016/j.infbeh.2014.05.009>
- Touchèque, M., Etienne, A.-M., Stassart, C., & Catale, C. (2016). Validation of the French version of The Parenting Stress Index-Short Form (fourth edition). *Journal of Community Psychology*, 44(4), 419-425. <https://doi.org/10.1002/jcop.21778>
- Vallotton, C., Harewood, T., Froyen, L., Brophy-Herb, H., & Ayoub, C. (2016). Child behavior problems: Mothers' and fathers' mental health matters today and tomorrow. *Early Childhood Research Quarterly*, 37, 81-93. <https://doi.org/10.1016/j.ecresq.2016.02.006>
- Van Steijn, D. J., Oerlemans, A. M., Van Aken, M. A. G., Buitelaar, J. K., & Rommelse, N. N. J. (2014). The reciprocal relationship of ASD, ADHD, depressive symptoms and stress in parents of children with ASD and/or ADHD. *Journal of Autism and Developmental Disorders*, 44(5), 1064-1076. <https://doi.org/10.1007/s10803-013-1958-9>
- Viladrich, C., Angulo-Brunet, A., & Doval, E. (2017). A journey around alpha and omega to estimate internal consistency reliability. *Anales de Psicología*, 33(3), 755. <https://doi.org/10.6018/analesps.33.3.268401>
- Whiteside-Mansell, L., Ayoub, C., McKelvey, L., Faldowski, R. A., Hart, A., & Shears, J. (2007). Parenting stress of low-income parents of toddlers and preschoolers: Psychometric properties of a Short Form of the Parenting Stress Index. *Parenting*, 7(1), 27-56. <https://doi.org/10.1080/15295190709336775>
- Williford, A. P., Calkins, S. D., & Keane, S. P. (2007). Predicting change in parenting stress across early childhood: Child and maternal factors. *Journal of Abnormal Child Psychology*, 35(2), 251-263. <https://doi.org/10.1007/s10802-006-9082-3>
- Zaidman-Zait, A., Miranda, P., Zumbo, B. D., Georgiades, S., Szatmari, P., Bryson, S., Fombonne, E., Roberts, W., Smith, I., Vaillancourt, T., Volden, J., Waddell, C., Zwaigenbaum, L., Duku, E., & Thompson, A. (2011). Factor analysis of the Parenting Stress Index-Short Form with parents of young children with autism spectrum disorders. *Autism Research*, 4(5), 336-346. <https://doi.org/10.1002/aur.213>